| DOCUMENT No.   DOCUMENT DESCRIPTION (Completed by Requestion Division)   Date of Request   Date of Req | UCN-7721B (2 1-86) MLB (2) 1-86) OAK RIDGE Y-12 PLANT INFORM                         | NATION CONTROL FORM  |             |
|--|--|--|-------------|
| Document Ne.  MS / ChR-003752 / K-25  Unclassified Titls: #   Storic   Chem   Cal   Kell   Cal   |  | rand By Requesting Division)   |             |
| Unclassified Titles: ### SECUTION CONTINUES AND CONTINUES  | Document No. Author's Telephone No.  | o. Acct. No. Date of Request   |             |
| Authorisi   SUTRAINE   Sandborg   Progress/Status Report   Co-Op Report   Thesu/Term Pacer   | MS/ChR-003752/K-25 $6-475$   | (125 - 0006) 1129193   |             |
| TYPE:   Formal Report   Informal Report   Progress/Status Report   Co-Op Report   Thass/Term Paper   | Unclassified Title: HIStoric Chemical K  | elease Report  | <del></del> |
| TYPE:   Formal Report   Informal Report   Progress/Status Report   Co-Op Report   Thass/Term Paper   |  |  |             |
| Oral Presentation (identify meeting, sponsor, location, data):   | Author(s) (Suzanne Sanabera)   |  |             |
| Journal Article (Identify Journal):     Other (Specify):     Occument will be published in proceedings   No   Yes  | TYPE: Formal Report Informal Report Progress/Statu                                   | us Report Co-Op Report Thesis/Term Paper   |             |
| Document will be published in proceedings   No   Yes   | Oral Presentation (identify meeting, sponsor, location, date):                       |  |             |
| Document will be published in proceedings   No   Yes   |  |  |             |
| Document will be published in proceedings  | Journal Article (Identify Journal):  |  |             |
| Document will be published in proceedings  | Other (Specify):   |  |             |
| Document will be distributed at meeting No Yes    No Yes   Yes   No Yes   N |  |  |             |
| Document has patent or invention significance   No   Yes (Identify)  |  |  |             |
| DOCUMENT has been previously released No Yes (Reference)    No   Yes (Reference)   | , -  |  |             |
| DIVISION REVIEW AND APPROVAL (Completed By Requesting Divisional TECHNICAL CLASSIFICATION REVIEW (Divisional Classification Representative) Title(s):   ACLOSSIFICATION REVIEW (Divisional Classification Representative)   DOCUMENT REQUEST APPROVED (Division or Department)   129/9 3   DOCUMENT: Lope   ACLOSSIFICATION CONTINUED STRIBUTION   DISTRIBUTION   DISTRIBUTION |  |  | -           |
| TECHNICAL CLASSIFICATION REVIEW (Divisional Classification Representative)  Title(s):   ACC Signature  |  |  |             |
| Title(s):   ACQSS Sect A Destract:   APPROVAL AND RELEASE    Case   Approval And Release subject to use of the following admonitory markings and conditions:   Destract Approval Copyright   Petent Caution   Petent Caution   Disclaimer   Copyright   Petent Caution   Disclaimer   Copyright   Petent Caution   Disclaimer   Copyright   Destract   Destract Approval Approved Approving admonitory markings and conditions:   Disclaimer   Copyright   Petent Caution   Other   Destract   Disclaimer   Copyright   Petent Caution   Other   Destract   Destra | • DIVISION REVIEW AND APPROVAL (C  | ompleted By Requesting Division)   |             |
| DOCUMENT: Legis   Abstract   Date   Date      Distribution Remarks:   Date   Date  |  | DOCUMENT REQUEST APPROVED (Division or Department)   | _           |
| Signature   Date   Da   |  | Jara 4 Welch 1/29/9  | <u> 3</u>   |
| Signature   Date   Da   | DOCUMENT: Legrel / Alla Sura Category  | Signature Date   |             |
| THE REMAINDER OF THIS FORM TO BE COMPLETED BY THE TECHNICAL INFORMATION OFFICE  DISTRIBUTION  Internal Distribution  THO File  Suzanne Sandberg  ANNOUNCE IN:  AND RELEASE  Date  APPROVAL AND RELEASE  Date  Date  Date  Category  Other  Date  Disclaimer   Copyright   Patent Caution   Other    Di | Java D. Welch 1/29/93  |  |             |
| Internal Distribution  | Signature Date   | Signature Date   |             |
| Internal Distribution   External Distribution   External Distribution   External Distribution   TiD-4500 Cetegory   or   Copies to OSTI   TiO File   Y-12 RC   Y-12    |  |  |             |
| External Distribution  |  |  |             |
| TID-4500 Category  |  | 1  |             |
| ANNOUNCE IN: ERA Atomindex (Available from NTIS)  M-3679 Category  ANNOUNCE IN: AWDR (Available from OSTI)  Distribution Remarks: AWDR (Available from OSTI)  APPROVAL AND RELEASE  Date Received  1-29-93  Date Initiated  |  | 1  |             |
| ANNOUNCE IN: AWDR (Available from OSTI) ANCR  Distribution Remarks: ALM RISK Clear Js. Public Black  APPROVAL AND RELEASE  Date Received 1-29-93   |  | Suzanne Sandberg   |             |
| Distribution Remarks: ALMANSK Cleans J. Pulse Black  APPROVAL AND RELEASE  Date Received 1-29-93   | M-3679 Category  |  |             |
| APPROVAL AND RELEASE  Date Received 1-29-93   Date Initiated 1-29-93    | ANNOUNCE IN: AWDR (Available from OSTI)  |  |             |
| APPROVAL AND RELEASE  Date Received 1-29-93   Date Initiated 1-29-93    | Chenific Chenif 2  | Police Polane  |             |
| Date Initiated  1-29-93  Date Initiated  1-29-93  Date  Date | Distribution Hemarks: GFFF NA / SA   | The Country of the Co |             |
| Date Initiated  1-29-93    Classifications:   Date   Date  |  |  |             |
| Title(s):   Molassfied   Abstract   Date   |  | RELEASE  |             |
| CLASSIFICATIONS:   |  |  |             |
| Title(s):  |  | Editor   |             |
| DOCUMENT:  Level McCassfud: Category   | Michael 3 J  | (// havin 1/29/9:  | <u>}</u>    |
| Level   Mclassfud   Category   Other   Date  | Abstract   | Stent Office / Date  |             |
| Weapons Data  Sigma  Other  Other  Date  V-12 Classification Office  NaOH data - Mandalay Chune (2 shorts)  APPROVED FOR:  Disclaimer Copyright  Patent Caution  Other  Technical Information Office  Date  Technical Information Office  Date  Date  Other  Technical Information Office  Date  Date  Other  Technical Information Office  Date   |  |  |             |
| Syd D  | Level (MClassificat Category   | Other Date   |             |
| V-12 Classification Office   Date     | Weapons Data Sigma   | Other Date   |             |
| A NaOH data - Mandalary / Churve (2 shorts)  Otherwise Confide dia 1 / R)  APPROVED FOR: Declassification Release subject to use of the following admonitory markings and conditions:  Disclaimer Copyright Patent Caution Other  Technical Information Office   | Lloyd & Firth 2/3/93   | Street   |             |
| ☐ Disclaimer ☐ Copyright ☐ Patent Caution ☐ Other ☐ Technical Information Office Dete  | * NaOH data - mandatary Remove (2 shorts)  |  |             |
| ☐ Disclaimer ☐ Copyright ☐ Patent Caution ☐ Other ☐ Technical Information Office Dete  | Otherwise Confidential IPD  APPROVED FOR: Declassification Release subject to use of | the following admonitory markings and conditions:  |             |
| Technical Information Office Dete  |  | P. VI - 7/1  | Sex         |
|  | பு பக்கோள் டு Copyright டி Patent Caution டி Other                                   | Technical Information Office Delo  | <del></del> |
|  | Conditions/Remarks:  |  |             |

## internal Correspondence

MARTIN MARIETTA ENERGY SYSTEMS, INC.

April 15, 1986

W. F. Furth

### Historic Chemical Release Report

Attached is the "Historic Chemical Release Report" for the Y-12 Plant. As can be noted, there are many gaps in the information. Some of this is because of the short time available but most of the gaps are because the information is not available. According to your instructions, we have provided the "best we could" in the time allowed. With more time, further information about the chemicals can be provided to complete the chemical release data form. Undoubtedly, a substantial effort could produce additional use information, but our efforts to date indicate the cost would far outweigh the value. An estimated 450 man-hours of the time of experienced and knowledgeable employees who are crucial to other work has already been spent in preparation of this report.

The list of chemicals selected was based on choices by several Y-12 individuals who have been involved in the use of chemicals during several years of plant operations. While it is no doubt arbitrary, we feel it represents a reasonable selection of chemicals of concern at Y-12 throughout the years. Materials not included are those in common public use such as fuels, cleaning supplies, or insecticides and those which may be hazardous but are used only in very small quantities and under very controlled conditions. A few other chemicals were selected for inclusion but no data was found concerning their usage.

The material provided is in many cases calculated or averaged since the records located did not provide year by year data. The extensive search made for usage and disposal records has revealed that in conversion from file to computer record systems, much of the historic data was eliminated, consistent with DOE instructions.

We feel we have provided essentially all the data available and will conclude our research with the submission of this report. Classified data forms will be forwarded under separate cover.

If you have questions, please contact W. G. Butturini at 4-3692.

BV Gordon G. Fee, 9704-2, MS-14 (4-2527) GGF:WGButturini:jll

RA Williams

Attachment: As Stated

cc/att.: W. G. Butturini - RC

D. T. Sissom

G. G. Fee

W. F. Thomas

W. L. Harper

J. E. Vath

G. E. Isham

L. F. Willis

M. L. Jones

|                             | Chemical Name: Ammonium Nitrate  |
|-----------------------------|--|
|                             | Uses:  |
|                             | Solid X Liquid Gas   |
|                             | Listed as Toxic: YesNo   |
|                             | Status of Environmental Compliance (Past and Present):   |
|                             |  |
| and the same                |  |
|                             | The state of the s |
| erepert to be send of the f | Known or Suspected Public Health Impacts:  |
|                             | May explode under confinement and high temperatures. On decomposition emits highly toxic fumes.  |
|                             | Known or Suspected Environmental Impacts:  |
|                             | Level and Type of Worker Protection:   |
|                             | Source of Chemical Usage Information:  |
|                             | Source of Chemical Distribution Information/Accuracy:  |

|                             | Unknown                  |                       |
|-----------------------------|--------------------------|-----------------------|
|                             | Other                    |                       |
|                             | Offsite<br>Burial        |                       |
| 34<br>=                     | Ons ite<br>Burial        |                       |
|                             |                          |                       |
|                             | Ons.                     |                       |
| Estimated Distribution in & | 1. Iquid<br>Effluent     | 100<br>* 000          |
| OT INVITED                  | Airborne<br>Release      | nitrate.              |
|                             | Product                  |                       |
|                             | Quantity<br>Used<br>(Kg) |                       |
|                             | Year                     | 1959<br>1958<br>mm su |
|                             | Chemical<br>Name         | charged as ammon      |

|                   | Chemical Name: Toluene Diisocyanate   |
|-------------------|---|
|                   | üses:   |
|                   | SolidLiquidX Gas  |
|                   | Listed as Toxic: Yes No   |
|                   | Status of Environmental Compliance (Past and Present):  |
|                   |   |
|                   |   |
| Pelo de accompany |   |
|                   | Known or Suspected Public Health Impacts:   |
|                   | Toxic by ingestion and inhalation, strong irritant to skin and tissues, especially eyes. Tolerance 0.02 ppm in air. |
|                   | Known or Suspected Environmental Impacts:   |
|                   | Level and Type of Worker Protection:  |
|                   | Source of Chemical Usage Information:   |
|                   | Source of Chemical Distribution Information/Accuracy  |

ESTIMATED CHEMICAL USAIL

Estimated Distribution in %

| Unknown                  |   |   |  |
|--------------------------|---|---|--|
| Other                    |   |   |  |
| Offsite<br>Burial        |   |   |  |
| Onsite<br>Burial         |   |   |  |
|                          |   | S   |  |
| Liquid Or<br>Effluent St |   | but no recon  |  |
| Airborne<br>Release      | 0.2<br>0.2<br>0.2<br>0.2<br>0.2                               | or to 1979  |  |
| Product                  | 002<br>002<br>002<br>002                                      | compound pr<br>-site.   |  |
| Quantity<br>Used<br>(Kq) | 11,045<br>8,386<br>9,409<br>8,795<br>8,782<br>7,364<br>10,023 | of this<br>buried or  |  |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980<br>1979          | ole use   |  |
| Chemical<br>Name         | Urethane foam<br>component<br>(toluene-<br>diisocyanate)      | There was considerable use of this compound pror to Considerable amounts were buried on-site. |  |

Estimated Distribution in X

| Unknown                  |   |
|--------------------------|---|
| 0ther                    |   |
| Offsite<br>Burial        |   |
| Ons i te<br>Buria l      |   |
| 00 S (                   |   |
| Liquid<br>Effluent       |   |
| Airborne<br>Release      | 84<br>87<br>96<br>94<br>85<br>99<br>100                                   |
| Product                  | ·   |
| Quantity<br>Used<br>(Kg) | 383,600<br>423,600<br>589,636<br>686,818<br>493,818<br>514,272<br>332,455 |
| Year                     | 1985<br>1984<br>1983<br>1980<br>1979                                      |
| Chemica 1<br>Namo        | Tetrachloroethylene   |

| Chemical Name: Chlorine   |
|---|
| Uses:   |
| SolidLiquidGas_X  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts: Chronic exposures can cause tooth corrosion, acne, and reduction in pulmonary function. It is a powerful skin, eye, and respiratory tract irritant and can cause fatal pulmonary edema.   |
| Known or Suspected Environmental Impacts:   |
| It is toxic to vegetation, white perch and striped bass, larvae, bluegill, minnows, and daphnia magna.  |
|   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation (2) rubber clothing (3) chemical goggles (4) face shield (5) proper respiratory protection. |
| Source of Chemical Usage Information:   |
|   |
| Same of Chamical Distribution Information/Accuracy:   |

L'STINVILLO CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |  |   |  |   |  |
|--------------------------|--|---|--|---|--|
| E                        |  |   |  |   |  |
| Other                    | 69*<br>41<br>30<br>27                  |   |  |   |  |
| Offsite<br>Burial        |  |   |  | · |  |
| Onsite<br>Burial         |  |   |  |   |  |
| G.                       |  |   |  |   |  |
| 2.2                      |  |   |  |   |  |
| 00                       |  |   |  |   |  |
| i. Iquld<br>Effluent     | 29***<br>57<br>68<br>71                | were removed  |  |   |  |
| Airborne<br>Release      | %<br>%<br>%<br>%<br>%                  | products w  |  |   |  |
| Product                  |  | The decomposition<br>um chlpride.                                       | oride.                                       |   |  |
| Quantity<br>Used<br>(Kg) | 27,000<br>177,000<br>158,500<br>58,000 | . The de  | odium chl                                    |   |  |
| Year                     |  | np oils<br>aimers.<br>to amm  | 0H to  |   |  |
| Chemica 1<br>Name        | Chlorine                               | *Trapped in vacuum pump oils. The decompos from the oils by reclaimers. | ***Neutralized with NaOH to sodium chloride. |   |  |

| Chemical Name: Copper Cyanide                          |
|--|
| Uses:  |
| Solid X Liquid Gas                                     |
| Listed as Toxic: YesNo                                 |
| Status of Environmental Compliance (Past and Present): |
|  |

Known or Suspected Public Health Impacts:

Following severe exposure to dust, symptoms of cyanide poisoning may develop. Ingest causes confusion, dizziness, unconsciousness, rapid pulse, convulsions, and paralysis Contact with eyes and skin causes irritation. Symptoms of inhalation exposure are similar to those caused by ingestion.

Known or Suspected Environmental Impacts:

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection.

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy

LSTINALLD CHEMICAL USAGE

|                    | Unknown                  | ·  |                            |  |      |
|--------------------|--------------------------|--|----------------------------|--|------|
|                    | Other                    | 100<br>100<br>100<br>100<br>100  |                            |  |      |
|                    | Offsite<br>Burial        | •  |                            |  |      |
| in L               | Onsite<br>Durial         |  |                            |  |      |
| 5                  | ere e per une            | THE CONTRACTOR OF THE PROPERTY |                            |  |      |
| stribut            | Onstite<br>Storage       | Section of the sectio | 700                        | the tile political design particular and | ·    |
| Estimated Distribu | Liquid<br>Effluent       |  |                            |  |      |
| ül                 | Airborne<br>Release      |  | of bath.                   |  |      |
|                    | Product                  |  | disposal                   |  |      |
|                    | Quantity<br>Used<br>(Kg) | 85<br>800<br>+350<br>275<br>923<br>967   | n prior to                 |  | ···· |
|                    | Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980<br>1979   | ing bat                    |  |      |
|                    | Chemical<br>Name         | Copper Cyanide   | *Destroyed in plating bath |  |      |

|         | Chemical Name: Tris (hydroxymethyl) nitromethane        |
|---------|---|
|         | Uses:   |
|         | Solid X Liquid Gas                                      |
|         | Listed as Toxic: Yes No                                 |
|         | Status of Environmental Compliance (Past and Present):  |
|         |   |
| -10 M/S |   |
|         |   |
|         | Known or Suspected Public Health Impacts:               |
|         | Moderate fire risk. Probably irritant to skin and eyes. |
|         |   |
|         | Known or Suspected Environmental Impacts:               |
|         |   |
|         |   |
|         |   |
|         | Level and Type of Worker Protection:                    |
|         |   |
|         |   |
|         | Source of Chemical Usage Information:                   |
|         |   |
|         |   |
|         | Source of Chemical Distribution Information/Accuracy:   |
|         |   |

LSTINALLD CHEMICAL USAGE

| 7 | 2                                     | -   |
|---|---------------------------------------|-----|
| - | =                                     |     |
|   | =                                     | *** |
|   | =                                     |     |
| 1 | _                                     |     |
| - |                                       |     |
|   |                                       |     |
| • | 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 |     |
|   | ٠                                     |     |
|   |                                       | - 1 |

| Unknown                  |                                  |   | · |   |  |
|--------------------------|----------------------------------|---|---|---|--|
| Other                    |                                  |   |   |   |  |
| Offsite<br>Burial        |                                  |   |   |   |  |
| Onsite<br>Burial         |                                  | Total Salah                                   |   |   |  |
| Onsite:                  | 1000%                            |   |   | • | The second of the second |
| L.Iquid<br>Effluent      |                                  | ed in West                                    | • |   |  |
| Airborne<br>Release      |                                  | to be treated                                 | · |   |  |
| Product                  | •                                | and stored                                    |   |   |  |
| Quantity<br>Used<br>(Kg) | 95<br>47.5<br>52.5<br>47.5       | facility                                      |   |   |  |
| Year                     | i                                | oolant<br>ity.                                |   |   |  |
| Themical Jame            | s (Hydroxy-<br>hyl)<br>romethane | *Treated at waste doolant facility and stored |   | • |  |

|           | Chemical Name: Zirconium Oxide   |
|-----------|--|
|           | Uses:  |
|           | Solid X Liquid Gas   |
|           | Listed as Toxic: YesNo   |
|           | Status of Environmental Compliance (Past and Present):   |
|           |  |
| A Section |  |
|           |  |
|           | Known or Suspected Public Health Impacts:  |
| •         | Tolerance 5 mg per cubic meter of air.   |
|           | · · · · · · · · · · · · · · · · · · ·  |
|           | Known or Suspected Environmental Impacts:  |
|           | WHOMIL OF GRabacter The second |
|           |  |
|           |  |
|           | Level and Type of Worker Protection:   |
|           | Level and Type of Worker Protection.   |
|           |  |
|           | To formation:  |
|           | Source of Chemical Usage Information:  |
|           |  |
|           | Source of Chemical Distribution Information/Accuracy:  |
|           |  |

LSTINALLD CHENICAL USAGE

|                        | Unknown                  |  |   |  |
|------------------------|--------------------------|--|---|--|
|                        | 0ther                    |  |   |  |
|                        | Offsite<br>Burial        |  |   |  |
| in &                   | Onsite<br>Burial         |  |   |  |
| 5                      |                          |  |   |  |
| stribu                 | Ons Ital                 | 100  |   |  |
| Estimated Distribution | Liquid<br>Effluent       | 100**<br>100<br>100<br>100<br>100            |   |  |
| 33                     | Airborne<br>Release      |  | in 1985.                                    |  |
|                        | Product                  | ·  | of waste discontinued in                    |  |
|                        | Quantity<br>Used<br>(Kg) | 1,398<br>1,091<br>818<br>727<br>590<br>318   | f waste d                                   |  |
|                        | Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | <del>-</del>                                |  |
|                        | Chemical                 | Zirconium Oxide                              | *As Raffinate.<br>**Direct liquid discharge |  |

|   | Chemical Name: Lead Carbonate                                    |
|---|--|
|   | Uses:  |
|   | Solid_X_LiquidGas  |
|   | Listed as Toxic: Yes   |
|   | Status of Environmental Compliance (Past and Present):           |
|   |  |
|   |  |
|   |  |
|   | Known or Suspected Public Health Impacts:                        |
|   | Toxic by inhalation. Tolerance - 0.15 mg per cubic meter of air. |
|   | Known or Suspected Environmental Impacts:                        |
|   |  |
|   |  |
|   | Level and Type of Worker Protection:                             |
| • |  |
|   | ·  |
|   | Source of Chemical Usage Information:                            |
|   |  |
|   | Source of Chemical Distribution Information/Accuracy:            |
|   | <del></del>  |

ESTINATED CHEMICAL USAGE

|                              | Unknown                  |                |                                   |  |        |   |              |
|------------------------------|--------------------------|----------------|-----------------------------------|--|--------|---|--------------|
|                              | 0ther                    |                |                                   |  |        |   |              |
|                              | Offsite<br>Burial        |                |                                   |  |        |   |              |
| in &                         | Onsite<br>Burial         |                |                                   |  | ······ |   | <br>         |
| strillition                  | Onsite<br>Storage        | *              | స్వాహిస్తున్నారు. క్రామాలు అయిది. | te a transport of the second seco |        | Maria e e e e e e e e e e e e e e e e e e e |              |
| Estimated Distribiltion in & | Liquid<br>Effluent       |                |                                   |  |        |   | <br>         |
| ات                           | Airborne<br>Release      |                |                                   |  | <br>   |   | <br><u>-</u> |
|                              | Product                  | 06             |                                   |  |        |   | <br>         |
|                              | Quantity<br>Used<br>(Kq) | 41,120         |                                   | <u>.</u>   | <br>   |   |              |
|                              | Year                     |                | •                                 |  | <br>   |   |              |
|                              | Chemical<br>Name         | Lead Carbonate |                                   | *Stored at K-25  |        |   |              |

| Chemical Name: Chromic Acid   |
|---|
| üses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: Yes No   |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Highly toxic, corrosive to skin. Tolerance 0.1 mg per cubic meter of air. Powerful oxidizing agent. May ignite on contact with organic materials. |
| Known or Suspected Environmental Impacts:   |
|   |
| _   |
| Level and Type of Worker Protection:  |
| Source of Chemical Usage Information:   |
| Source of Chemical Distribution Information/Accuracy:   |

ESTINATED CHEMICAL USAGE

|                           | Unknown                  |  |
|---------------------------|--------------------------|--|
|                           | Other .                  | <i>₹</i>                                 |
|                           | Offsite<br>Burial        |  |
| in &                      | Onsite<br>Burial         |  |
| Estimated Distribution in | Onsalte<br>Storane       | 000000<br>000000                         |
| timated Di                | Liquid<br>Effluent       |  |
| ŭ)                        | Airborne<br>Release      |  |
|                           | Product                  |  |
| -                         | Quantity<br>Used<br>(Kg) | 241<br>265<br>275<br>214<br>1,270<br>649 |
|                           | Year                     |  |
|                           | Chemical<br>Name         | Chromic Acid                             |

| C        | hemical Name: Furfuryl Alcohol  |
|----------|---|
| Us       | ses:  |
| So       | olidLiquidXGas  |
| L        | isted as Toxic: Yes   |
| S1       | tatus of Environmental Compliance (Past and Present):   |
|          |   |
|          |   |
| X:       | nown or Suspected Public Health Impacts:  |
| To       | oxic by inhalation. May react explosively with mineral nd some organic acids. Tolerance - 5 ppm in air. |
| K.       | nown or Suspected Environmental Impacts:  |
| <u>r</u> | evel and Type of Worker Protection:   |
| s        | Source of Chemical Usage Information:   |
| <br>s    | Source of Chemical Distribution Information/Accuracy:   |

ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |                      |   |
|--------------------------|--|----------------------|---|
| 0ther                    | 404<br>400<br>400<br>400<br>400<br>400<br>400<br>400<br>400<br>400                                 |                      |   |
| Offsite<br>Burial        |  |                      |   |
| Onsite<br>Burial         |  |                      |   |
| Onstre<br>Storage        | · <u>Lucia e rei</u> fficación.  |                      |   |
| Liquid<br>Effluent       |  |                      |   |
| Airborne<br>Release      |  |                      | • |
| Product                  | 090000000000000000000000000000000000000  | site.                |   |
| Quantity<br>Used<br>(Kg) | 4,580<br>4,580<br>5,725<br>9,160<br>11,450<br>11,450<br>11,450<br>6,870<br>6,870<br>6,870<br>4,580 | some off             |   |
| Year                     | 1973<br>1972<br>1971<br>1970<br>1969<br>1966<br>1966<br>1965<br>1965<br>1961                       | site,                | · |
| Chemical<br>Name         | Furfuryl Alcohol<br>Polymer  | *Incinerated some on |   |

| Chemical Name: Potassium Cyanide  |
|---|
| üses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
| Known or Suspected Public Health Impacts:  Symptoms which can occur from either oral or inhalation exposure are sore throat, faintness, labored breathing and abdominal pain. Severe cases can result in convulsions and death. Contact with skin can cause chemical burns and with eyes, blurred vision and burns. |
| Known or Suspected Environmental Impacts:   |
| ·   |
| ·   |
|   |

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection.

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy:

LSTINALED CHEMICAL HSAGE

Estimated Distribution in &

|                  | Unknown  |  |                       | <br> |      |                   |
|------------------|----------|--|-----------------------|------|------|-------------------|
|                  | Other    | 100%*<br>100%<br>100%<br>-<br>100%<br>100%   | <br>                  |      |      |                   |
| Offsite          | Burial   |  |                       | <br> |      |                   |
| Onsite           |          |  |                       |      | <br> | <del></del> -     |
| Ons ite          | Storage  |  |                       |      | <br> |                   |
| Liguid           | Effluent |  |                       |      |      |                   |
| _<br>Airborne    | Release  |  | of bath.              | <br> |      | <del>.     </del> |
|                  | Product  |  |                       |      | <br> |                   |
| Quantity<br>Used | (Kg)     | 1<br>1<br>3<br>3<br>3                        | bath prior to disposa |      |      |                   |
|                  | Year     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 |                       | <br> |      | ·                 |
| Chemical.        | Name     | Potassium Cyanide                            | *Destroyed in plating |      |      |                   |

| Chemical Name: Perchloric Acid   |
|--|
| Uses:  |
| SolidLiquidX Gas   |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
| Known or Suspected Public Health Impacts:  Acute exposure by inhalation causes a burning sensation in nose and throat and orally, blistering and chemical burns to mouth, esophagus, and stomach. Chemical burns of eyes and skin occur following acute exposure. Vomiting and severe coughing can result from prolonged or excessive exposure.  Known or Suspected Environmental Impacts: |
| Level and Type of Worker Protection:  Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation (2) rubber clothing (3) chemical goggles (4) face shield (5) proper respiratory protection.   |

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy:

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |  |  |
|--------------------------|--|--|--|
| 0ther                    |  |  |  |
| Offsite<br>Burial        |  |  |  |
| Onsite<br>Burial         |  |  |  |
| Ons te<br>Storage        | **06<br>06<br>06<br>06<br>06<br>06           | ic acid as   |  |
| Liquid<br>Effluent       |  | jas and the hydrochloric<br>newhat larger.   |  |
| Airborne<br>Release      | 10%<br>10%<br>10%<br>10%<br>10%              |  |  |
| Product                  |  | ease of the<br>probably sc   |  |
| Quantity<br>Used<br>(Kg) | 429<br>384<br>290<br>334<br>111<br>50        | acid.<br>n the rel   |  |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | nloric<br>ed bot<br>Quanti   |  |
| Chemical                 | Perchloric Acid                              | *As HCl gas.  **Primarily as hydrochloric acid.  Earlier years included both the release of a liquid effluent. Quantities were probabl |  |

| Chemical Na    | me: Sodium Cyanide                           |
|----------------|--|
| üses:          |  |
| Solid <u>X</u> | Liquid X Gas                                 |
|                | Coxic: Yes                                   |
| Status of E    | Invironmental Compliance (Past and Present): |
|                |  |

Known or Suspected Public Health Impacts:

It may be absorbed into the body by inhalation, ingestion, and through the skin. Symptoms which can result from either oral or inhalation exposure are sore throat, faintness, labored breathing and abdominal pain. Serious cases may result in convulsions and death. Contact with skin can cause burns and with eyes, blurred vision and chemical burns.

Known or Suspected Environmental Impacts:

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection.

Source of Chemical Usage Information:

Source of Chemical Distribution: Information/Accuracy:

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Other Unknown            | 100%*<br>100%<br>100%<br>100%<br>100%        |              |            |       |   |  |
|--------------------------|--|--------------|------------|-------|---|--|
| Offsite<br>Burial        |  |              |            | <br>  | • |  |
| Onsite<br>Burial         |  |              |            | <br>  |   |  |
| Onsite<br>Storage        |  |              |            | <br>  |   |  |
| Liquid<br>Effluent       |  |              |            | <br>  |   |  |
| Airborne<br>Release      |  | before 1979. |            |       |   |  |
| Product                  |  | years        |            |       |   |  |
| Quantity<br>Used<br>(Kg) | 91<br>-<br>1,000<br>363<br>273<br>954<br>956 | number of    |            | <br>- |   |  |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | And a        |            |       |   |  |
| Chemica I<br>Name        | Sodium Cyanide                               | ·            | *Destroyed |       |   |  |

| Chemical Name: Chloroform   |
|---|
| Uses:   |
| SolidLiquidXGas   |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts: Chronic exposure can damage liver and kidney. Acute inhalation exposure can cause CNS depression, GI disturbances, intracranial pressure changes, respiratory disorders, and heart arrest. Acute oral intake can cause GI disturbances, unconsciousness and possibly liver and kidney damage. Acute skin and eye contact causes pain and severe irritation of eyes. |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection: Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respiratory protection.  |
| Source of Chemical Usage Information:   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |
| SOUTCE OF CHempony Productions  |

ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |
|--------------------------|--|
| Other Other              |  |
| Offsite<br>Burial        |  |
| Onsite<br>Burial         |  |
| Onsite<br>Storage        | 006<br>066<br>066                            |
| Liquid<br>Effluent       |  |
| Airborne<br>Release      | 010<br>010<br>10<br>10<br>10<br>10<br>10     |
| Product                  |  |
| Quantity<br>Used<br>(Kg) | 214<br>158<br>203<br>169<br>174<br>73<br>336 |
| Year                     | 1985<br>1984<br>1983<br>1980<br>1979         |
| Chemica)<br>Name         | Chloroform                                   |

|   | Chemical Name: Methyl Ethyl Ketone   |
|---|--|
|   | Uses:  |
|   | SolidLiquidX Gas   |
|   | Listed as Toxic: YesNo   |
|   | Status of Environmental Compliance (Past and Present):   |
|   |  |
|   |  |
|   |  |
|   | Known or Suspected Public Health Impacts: Chronic exposure can cause headar throat irritation, dermatitis, and numbness of fingers (affects peripheral nervous system). Acute inhalation exposure causes nose and throat irritation, headache, and dizziness. CNS depression may occur depending on concentration. Acute oral exposure may cause vomiting and nausea. It is skin and eye irritant. |
|   | Known or Suspected Environmental Impacts:  |
|   |  |
|   |  |
|   |  |
|   | Level and Type of Worker Protection:   |
|   | Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respiratory protection.   |
|   | Source of Chemical Usage Information:  |
|   |  |
| • | Toformation / locuracy:  |
|   | Source of Chemical Distribution Information/Accuracy:  |

# ESTINATED CHEMICAL HSAGE

Estimated Distribution in %

| Unknown                  |   |  |  |
|--------------------------|---|--|--|
| 0ther                    |   |  |  |
| Offsite<br>Burial        |   |  |  |
| Onsite<br>Burial         |   |  |  |
| Onsite<br>Storage        | 88<br>75<br>75<br>75<br>75                              | ia].   |  |
| Liquid<br>Effluent       |   | solvent/cleaning material<br>fluent.   |  |
| Airborne<br>Release      | 12<br>25<br>25<br>25<br>25<br>25<br>25                  | as solvent/cle<br>effluent.  |  |
| Product                  |   | as<br>eff  |  |
| Quantity<br>Used<br>(Kg) | 495.4<br>678.6<br>576.8<br>740.3<br>692<br>500<br>1,005 | ears in q<br>ne and in   |  |
| Year                     |   | lier y<br>airbor   |  |
| Chemica l<br>Name        | yl Ethyl Ketone   | MEK also used in earlier years in quantity<br>Significant amounts airborne and in liquid |  |

| Chemical | Name: Benze | ene             |       |     |           |  |
|----------|-------------|-----------------|-------|-----|-----------|--|
| Uses:    |             |                 |       |     |           |  |
| Solid    | Liquid_     | X Gas           | ·     |     |           |  |
| Listed a | s Toxic: Ye | s No            |       |     |           |  |
| Status c | f Environme | ntal Compliance | (Past | and | Present): |  |

Known or Suspected Public Health Impacts: Chronic exposure may cause blood disorders, including aplastic anemia and leukemia, embryonal death and fetotoxicity and chromosomal aberrations (bone marrow and lymphocytes). Acute effects from ingestion or inhalation are irritation of mucous membranes, convulsions, CNS depressi and respiratory failure resulting in possible death. Harmful amounts may be absorbed through skin. It is an irritant to the skin and eyes and causes transient injury to Known or Suspected Environmental Impacts: cornea. It causes dry skin.

Biological effects reported in algae, protozoa, grass shrimp, crab larvae, shrimp, bluegill, fathead minnows, bass, herring and anchovy larvae, brown trout and clawed toads.

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respirate protection.

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy:

L'STIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |   |                       |            |             |
|--------------------------|---|-----------------------|------------|-------------|
| Other                    |   |                       |            |             |
| Offsite<br>Burial        |   |                       |            |             |
| Onsite<br>Burial         |   |                       |            |             |
| Onsite<br>Storage        | 8<br>8<br>8<br>8<br>8<br>8<br>8<br>8<br>8 |                       |            |             |
| Liquid<br>Effluent       |   |                       |            |             |
| Airborne<br>Release      | 15<br>15<br>15<br>15                      |                       |            | <del></del> |
| Product                  |   |                       |            |             |
| Quantity<br>Used<br>(Kg) | 35<br>31<br>59<br>41                      | estimates.            | <u>-</u> - |             |
| Year                     |   | are a                 |            |             |
| Chemical'<br>Name        | Benzene                                   | % distributions above | •.         |             |

| Chemica | ıl N | lame: | Toule | ne   |            |       |     |           |
|---------|------|-------|-------|------|------------|-------|-----|-----------|
| üses:   |      |       |       |      |            |       |     |           |
| Solid_  | _    | _ Liq | uid_  | Х    | Gas        | ·     |     |           |
| Listed  | as   | Toxic | : Yes | ·    | . Но       | ,     |     |           |
| Status  | of   | Envir | onmer | ital | Compliance | (Past | and | Present): |

Known or Suspected Public Health Impacts: Chronic effects are defatting of skin and hematologic disorders. Brain injury may result from chronic sniffing. It is an eye and respiratory irritant. High vapor concentration causes drowsiness, vision disorders, and possibly cardiac arrhythmias and coma. If aspirated it may cause pulmonary edema. Orally it can cause nausea, vomiting, abdominal pain, diarrhea, and respiratory depression. Liver, kidney, and heart injury are possible.

Known or Suspected Environmental Impacts:

It can be lethal to rainbow trout, channel catfish and bluegill.

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respirator protection.

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy:

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |  |          |   |
|--------------------------|--|--|----------|---|
| Other                    |  |  |          |   |
| Offsite<br>Burial        |  |  | ,        |   |
| Onsite<br>Burial         |  |  |          |   |
| Onsite<br>Storage        | 066688 <u>8</u>                                  | no<br>air.   |          |   |
| Liquid<br>Effluent       |  | earlier years but there are<br>rom plant and released into                             |          |   |
| Airborne<br>Release      | 10<br>10<br>10<br>10<br>20<br>20                 | er years b<br>lant and re  |          | · |
| Product                  |  | ies in earl<br>ents from p   |          |   |
| Quantity<br>Used<br>(Kg) | 161<br>393<br>268<br>251<br>367<br>311           | r quantit<br>uid efflu   | <u>-</u> |   |
| Year                     | <br>1985<br>1984<br>1983<br>1982<br>1980<br>1979 | greate<br>the life   |          |   |
| Chemical<br>Name         | Toluene  | Toluene used in much greater quantities in records. It was in the liquid effluents for |          |   |

| Chemica | 1 : | Name:   | Hydroflud | oric Acid  |       |     |           |
|---------|-----|---------|-----------|------------|-------|-----|-----------|
| Uses:   |     |         |           |            |       |     |           |
| Solid   |     | Liqui   | d X       | Gas        | ·     |     |           |
| Listed  | as  | Toxic:  | Yes       | No         |       |     |           |
| Status  | οî  | Environ | nmental   | Compliance | (Past | and | Present): |
|         |     |         |           |            |       |     |           |

Known or Suspected Public Health Impacts:

Chronic effects which may result from inhalation or ingestion are fluorosis weight loss, malaise, anemia, leukopenia, discoloration of teeth and osteosclerosis. Repeated exposure to low concentration of gas may cause nasal congestion, bronchitis, bone changes and chronic irritation of nose and throat. Acute exposure to skin causes burns and ulceration. Liquid or vapor causes chemical burns and permanent visu Known or Suspected Environmental Impacts: defects or blindness. More sever effects may occur depending on concentration.

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include:

(1) adequate ventilation (2) rubber clothing (3) chemical goggles (4) face shield

(5) proper respiratory protection.

Source of Chemical Usage Information:

Source of Chemical Distribution Information/Accuracy:

### ESTIMATED CHEMICAL USAGE

Estimated Distribution in 1

| •   |  | ٠                                       |          |   |   |                   |                  |                   |       |         |
|---|--|---|----------|---|---|-------------------|------------------|-------------------|-------|---------|
| Chemical                                  | Year   | Quantity<br>Used                        | Provinct | Airborne<br>Release                           | l.Iquid<br>Effluent   | Onsite<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other | Unknown |
| rofluoric Acid<br>ntinued)                | 1958<br>1957<br>1956<br>1955<br>1954<br>1953<br>1951 | 000000000000000000000000000000000000000 |          | .05<br>.05<br>.05<br>.05<br>.05<br>.05<br>.05 | 67.95<br>67.95<br>67.95<br>24.95<br>24.95<br>24.95<br>24.95 |                   |                  | •                 |       |         |
|   | 1946<br>1945   | 4,000<br>2,600                          | 300      |   | 70*   |                   |                  |                   |       |         |
| *Discharged in caustic scrubber solution. | tic scr  | ubber sol                               | ution.   |   |   |                   |                  |                   |       |         |

### LSTIMALLD CHEMICAL USAGE

# Estimated Distribution in X

| Unknown                  |  |
|--------------------------|--|
| Other                    |  |
| Offsite<br>Burial        | •  |
| Onsite<br>Durial         |  |
| Onstte<br>Storage        |  |
| Liquid<br>Effluent       | 25<br>25<br>25<br>25<br>25<br>25<br>25<br>24<br>25<br>24<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25<br>25   |
| Airborne<br>Release      |  |
| Product                  | <b>755 755</b> |
| Quantity<br>Used<br>(Kg) | 5,000<br>5,000<br>5,000<br>5,000<br>5,000<br>5,000<br>11,000<br>11,000<br>11,000   |
| Year                     | 1984<br>1984<br>1983<br>1983<br>1981<br>1975<br>1977<br>1977<br>1977<br>1977<br>1967<br>1967<br>1967<br>1967   |
| Shemical Sheme           | Hydrofluoric Acid  |

| Chemical Name: Aluminum Nitrate                        |
|--|
| Uses:  |
| Solid X Liquid Gas                                     |
| Listed as Toxic: Yes No                                |
| Status of Environmental Compliance (Past and Present): |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:              |
| Strong oxidizing agent.                                |
|  |
| Known or Suspected Environmental Impacts:              |
|  |
|  |
|  |
| Level and Type of Worker Protection:                   |
|  |
|  |
| Source of Chemical Usage Information:                  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |

### I STINATED CHEMICAL USAGE

### BETA PROCESS

Estimated Distribution in %

| Unknown                  |                              |                                  |  |      |      |
|--------------------------|------------------------------|----------------------------------|--|------|------|
| Other .                  |                              |                                  |  |      |      |
| Offsite<br>Burial        |                              |                                  |  |      | <br> |
| Ons Ite<br>Burial        |                              |                                  |  |      | <br> |
| Ons I te<br>Storage      |                              |                                  |  | <br> |      |
| L. Iquid<br>Effluent     | 100*                         |                                  |  |      |      |
| Airborne<br>Release      |                              | solution.                        |  |      |      |
| Product                  |                              | aluminum fluoride in a nitrate s |  | <br> |      |
| Quantity<br>Used<br>(Kg) | 16,500<br>11,000             | uoride in                        |  |      |      |
| Year                     | <del> </del><br>1946<br>1945 | num fl                           |  |      |      |
| Chemica 1<br>Name        | Aluminum Nitrate             | *Discharged as alumi             |  |      |      |

| Chemical Name: Ammonium Hydroxide   |
|---|
| Uses:   |
| SolidLiquid_X_Gas   |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Chronic exposure causes respiratory disorders. Acute inhalation exposure causes severe respiratory irritation resulting in pulmonary edema, bronchitis, respiration disorders, vomiting and nausea. Acute oral intake causes corrosion of upper GI tract and chemical burns. Acute exposure of eyes and skin results in irritation and chemical burns.  Known or Suspected Environmental Impacts: |
| Level and Type of Worker Protection:  Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation (2) rubber clothing (3) chemical goggles (4) face shield (5) proper respiratory protection.   |
| Source of Chemical Usage Information:   |
|   |
|   |

Source of Chemical Distribution Information/Accuracy:

LSTINALLD CHEMICAL USAGE

Estimated Distribution in &

| Chemica 1          | Year              | Quantity<br>Used<br>(Kg) | Product | Airborne<br>Release | l.Iquid<br>Effluent | Ons I te<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other . | Unknown |
|--------------------|-------------------|--------------------------|---------|---------------------|---------------------|---------------------|------------------|-------------------|---------|---------|
| Ammonium Hydroxide | 1980              | 75,000                   | -       |                     | *66<br>-            |                     |                  |                   |         |         |
| 1-7- E-EV          | 1979              | 75,000                   |         |                     | *66                 |                     |                  |                   |         |         |
|                    | 1977              | 75,000                   |         |                     | *66                 |                     |                  |                   |         |         |
| -                  | 1975              | 75,000                   |         |                     | *66                 |                     |                  |                   |         |         |
|                    | 1974              | 75,000                   |         |                     | * *66               |                     |                  |                   |         |         |
|                    | 1973              | 75,000                   |         |                     | *66                 |                     |                  |                   |         |         |
| -<br>-             | 1972              | 75,000                   | -       |                     | *66<br>-            |                     |                  |                   |         |         |
|                    | 1970              | 40,000                   |         |                     | *66                 |                     |                  |                   |         |         |
|                    | 1968              | 40,000                   | •       | ← -                 | *66                 |                     |                  |                   |         |         |
|                    | 1967<br>1966      | 40,000                   |         |                     | *66<br> -           |                     |                  |                   |         |         |
| . Y 6              | 1965              | 40,000                   |         | <b>.</b>            | *66                 |                     |                  |                   |         |         |
|                    | 1964<br>1963      | 40,000                   |         |                     | *66<br>             |                     |                  |                   |         |         |
|                    | 1962              | 40,000                   |         | , <del>, , ,</del>  | *66                 |                     |                  |                   |         |         |
|                    | 1961              | 40,000                   |         |                     | *66<br>             |                     |                  |                   |         |         |
|                    | 1959              | 55,000                   |         |                     | **66                |                     |                  |                   |         |         |
|                    | 1958              | 55,000                   |         | <b>.</b>            | **66                |                     |                  |                   |         |         |
|                    | 1957              | 75,000                   |         | <u>-</u>            | **66                |                     |                  | ***               |         |         |
|                    | 1956              | 75,000                   |         | <b></b> -           | **66<br>            |                     |                  | •                 |         |         |
| as                 | ammonium nitrate. | trate.                   | inomae  | m fluoride          |                     |                     |                  |                   |         |         |
|                    |                   | רו מרכ                   |         | 2                   |                     |                     |                  |                   |         |         |
|                    |                   |                          |         | <del>.    </del>    |                     |                     |                  |                   |         |         |

LSTINALLD CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |   |                              |  |  |
|--------------------------|---|------------------------------|--|--|
| 0ther                    |   |                              |  |  |
| Offsite<br>Burial        |   |                              |  |  |
| Onsite<br>Ourial         |   |                              |  |  |
| Onsite<br>Storage        |   |                              | fate.  |  |
| L. Iquid<br>Effluent     | ****66<br>****66<br>***66                           | ****66<br>***66              | ammonium sulfate.  |  |
| Airborne<br>Release      |   |                              | m fluoride.  |  |
| Product                  |   |                              | ammoniu  |  |
| Quantity<br>Used<br>(Kg) | 195,000<br>195,000<br>285,000<br>855,000<br>815,000 | 315,000<br>230,000<br>85,000 | itrate.<br>itrate an<br>nitrate,                                     |  |
| Year                     | 1955<br>1954<br>1953<br>1952<br>1951<br>1950        | 1945<br>1944<br>1943         | mmenium r<br>ammonium r  |  |
| Chemical<br>Name         | Ammonium Hydroxide<br>(continued)                   |                              | *Discharged as ammo<br>**Discharged as ammo<br>***Discharged as ammo |  |

| Chemical Name: Phthalic Acid   |
|--|
| üses:  |
| Solid X Liquid Gas   |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):               |
|  |
|  |
| •  |
| Known or Suspected Public Health Impacts:                            |
| Irritating to mucuous membranes and in high concentrations narcotic. |
| Known or Suspected Environmental Impacts:                            |
|  |
| Level and Type of Worker Protection:                                 |
| Source of Chemical Usage Information:                                |
| Source of Chemical Distribution Information/Accuracy                 |

### L'STINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                   |                       |   |      | <br> |  |
|--------------------------|-------------------|-----------------------|---|------|------|--|
| Other .                  |                   |                       |   |      | <br> |  |
| Offsite<br>Burial        |                   |                       |   |      |      |  |
| Onsite<br>Burial         |                   |                       |   | <br> |      |  |
| Onsite<br>Storage        |                   |                       |   | <br> |      |  |
| Liquid<br>Effluent       | 100*<br>100       |                       |   |      |      |  |
| Airborne<br>Release      |                   |                       |   |      |      |  |
| Product                  |                   |                       |   |      |      |  |
| Quantity<br>Used<br>(Kg) | 13,000<br>115,000 | nalate.               |   |      |      |  |
| Year                     | ļ<br>1953<br>1952 | ammonium phthalate.   |   |      |      |  |
| Chemica 1<br>Name        | Acid              | *Discharged as ammoni | ÷ |      |      |  |

| Chemical Name: Salicylic Acid   |
|---|
| Uses:   |
| Solid_XLiquidGas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
| •   |
| Known or Suspected Public Health Impacts:   |
| Ingestion of large amounts can cause vomiting, abdominal pains, increased respiration, acidosis, and mental disturbances. May cause rashes. |
| Known or Suspected Environmental Impacts:   |
|   |
| Level and Type of Worker Protection:  |
| Source of Chemical Usage Information:   |
| Source of Chemical Distribution Information/Accuracy:   |

L'STINATED CHEMICAL USAGE

Estimated Distribution in X

| J.                       |                   |   |                         |      |      |
|--------------------------|-------------------|---|-------------------------|------|------|
| Unknown                  |                   |   |                         |      | <br> |
| Other .                  | 95**              |   |                         |      |      |
| Offsite<br>Burial        |                   |   |                         |      |      |
| Ons i te<br>Aur ia 1     |                   |   |                         | <br> |      |
| Onsite<br>Storage        |                   |   |                         | •    |      |
| l. Iquid<br>Effluant     | <br>              |   | •                       |      |      |
| Airborne<br>Release      |                   |   |                         |      |      |
| Product                  |                   |   | solution.               |      |      |
| Quantity<br>Used<br>(Kg) | 700,000<br>78,000 | - | chloride s              |      | <br> |
| Year                     | i<br>1951<br>1952 |   | nium o<br>ysis.         | <br> | *    |
| Chemica I<br>Name        | Salicylic Acid    |   | *Discharged in ammonium |      |      |

| Chemical Name: Zirconium Tetrachloride   |
|--|
| Uses:  |
| Solid X Liquid Gas   |
| Listed as Toxic: YesNo   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Toxic and irritant. Tolerance - 5 mg per cubic meter in air.   |
|  |
| Known or Suspected Environmental Impacts:  |
|  |
| ·  |
|  |
| Level and Type of Worker Protection:   |
| •<br>·   |
|  |
| Source of Chemical Usage Information:  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
| TOUTGE OF CHEMPORE PERCENTIAL PROFESSION AND ASSESSION AND ASSESSION AND ASSESSION AND ASSESSION AND ASSESSION ASSES |

LSTIMATED CHEMICAL USAGE

Estimated Distribution in 2

| Unknown                  | .6 as hafnium                          |   |      |   |  |
|--------------------------|--|---|------|---|--|
| 0ther                    |  |   |      |   |  |
| Offsite<br>Burial        |  |   |      |   |  |
| Onsite<br>Burial         |  |   |      |   |  |
| Onsite<br>Storage        |  |   | <br> |   |  |
| l.Iquid<br>Effluent      | .02 .02 .02 .02                        |   |      |   |  |
| Airborne<br>Release      | .02<br>.02<br>.02                      |   |      |   |  |
| Product                  |  |   | <br> |   |  |
| Quantity<br>Used<br>(Kg) | 72,000<br>649,000<br>648,000<br>72,000 |   |      |   |  |
| Year                     |  |   |      |   |  |
| Chemica)                 |  | · |      | *************************************** |  |

| Chemical Name: Methyl Isobutyl Ketone                       |
|---|
| Uses: Solvent   |
| SolidLiquid_XGas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):      |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:                   |
| Flammable, dangerous fire risk. Tolerance - 100 ppm in air. |
|   |
| Known or Suspected Environmental Impacts:                   |
|   |
|   |
|   |
| Level and Type of Worker Protection:                        |
|   |
|   |
| Source of Chemical Usage Information:                       |
|   |
| Source of Chemical Distribution Information/Accuracy:       |
| Source of Chemical Distribution information/ scource.       |

LSTINALLD CHEMICAL USAGE

Estimated Distribution in X

| Unknown                  |  |  |  |
|--------------------------|--|--|--|
| 0ther                    | 30<br>30<br>30<br>30                   |  |  |
| Offsite<br>Burial        |  |  |  |
| Onsite<br>Burial         |  |  |  |
| Ons I te<br>Storage      |  | tc.  |  |
| l. Iquid<br>Effluent     | *09<br>09                              | ammonium sulfate,  |  |
| Airborne<br>Release      | 10<br>10<br>10                         | oride, ammoniu   |  |
| Product                  |  | ammonium chlori<br>calcination.                                |  |
| Quantity<br>Used<br>(Kg) | 16,000<br>136,000<br>136,000<br>15,000 | ı <u>B</u>   |  |
| Year                     | 1953<br>1952<br>1951<br>1950           | e stre<br>ysis d   |  |
| Chemical<br>Name         | Hexone<br>(Methyl Isobutyl<br>Ketone)  | *Discharged as waste streams<br>**Destroyed by pyrolysis durir |  |

| Chemical Name: Hafnium   |
|--|
| üses:  |
| Solid X Liquid Gas   |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Explosive in powder form, either dry or wet with less than 25% water. Tolerance - 0.5 mg per cubic meter of air. |
| Toterance - 0.5 mg per cubic meses of and  |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
| •  |
| Level and Type of Worker Protection:   |
|  |
|  |
| Source of Chemical Usage Information:  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

LSTINALLD CHEMICAL USAGL

Estimated Distribution in %

|              |                              | Quantity<br>Used             | Descript             | Airborne<br>Release | t. iquid<br>Effluent | Onsite<br>Storage | Onsite<br>Aurial | Offsite<br>Burial | 0ther | Unknown |
|--------------|------------------------------|------------------------------|----------------------|---------------------|----------------------|-------------------|------------------|-------------------|-------|---------|
|              | 1953<br>1952<br>1951<br>1950 | 500<br>4,000<br>4,000<br>500 | 50<br>50<br>50<br>50 |                     | 50*<br>50<br>50      |                   |                  | •                 |       |         |
| This element | was s                        | separated <sup>F</sup>       | rom zirconium.       | ·wr                 |                      |                   |                  |                   |       |         |
|              | m hy                         | as hafnium hydroxide so      | solution.            |                     |                      |                   |                  |                   |       |         |
|              |                              |                              |                      |                     |                      |                   |                  |                   |       |         |
|              |                              |                              |                      |                     |                      |                   |                  |                   |       |         |
|              |                              |                              |                      |                     | <del></del>          |                   |                  |                   |       |         |

| Chemical Name: Asbestos   |
|---|
| Uses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| It causes delayed toxicity to respiratory tract seen after long periods of exposure. It may cause lung cancer, dyspnea, dry cough, reduced appetite, and weakness.  |
| Known or Suspected Environmental Impacts:   |
|   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection, paper suits. |
| Source of Chemical Usage Information:   |
| Source of Chemical Distribution Information/Accuracy:   |

LSTINALLD CHEMICAL USAGE

Estimated Distribution in 8

| Unknown                  | ,   |                      |   |
|--------------------------|---|----------------------|---|
| Other .                  |   |                      |   |
| Offsite<br>Burial        | ·   |                      |   |
| KGS<br>Onsite<br>Burial  | 168,794<br>50,018<br>72,270<br>45,482<br>18,580<br>20,324<br>36,762<br>6,425* |                      |   |
| Ons I te<br>Storage      |   |                      |   |
| L.Iquid<br>Effluent      | <del>-</del>  |                      |   |
| Airborne<br>Release      | ·   |                      |   |
| Product                  |   | in 1978.             |   |
| Quantity<br>Used<br>(Kg) |   | osal begar           | · |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1979<br>1978                          | s disposal           | • |
| Chemical<br>Name         | Asbestos-Bearing<br>Materials   | *Records of asbestos |   |

| Chemical Name: Sodium Dichromate  |
|---|
| üses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
| •   |
| Known or Suspected Public Health Impacts: Liver and CNS damage can result from chronic exposure. Acute inhalation exposure causes irritation, shortness of breath, chest pain, and chemical pneumonia. Acute oral exposure causes corrosion of mucous membranes, gastroenteritis, vascular collapse delirium, liver damage and renal failure. It is corrosive to skin and eyes. |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
| •   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material incl (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection.  |
| Source of Chemical Usage Information:   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |

## LSTINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |  |     | <br>- <del></del> |  |
|--------------------------|--|--|-----|-------------------|--|
| Other                    |  |  |     |                   |  |
| Offsite<br>Burial        |  |  |     |                   |  |
| Onsite<br>Aurial         |  |  |     | <br>              |  |
| Onsite<br>Storage        |  |  |     |                   |  |
| l. Iquid<br>Effluent     |  |  |     |                   |  |
| Airborne<br>Ne lease     |  | to estimate.<br>nts and is<br>rds prior  |     |                   |  |
| Product                  |  | ~ ~  |     | <br>              |  |
| Quantity<br>Used<br>(Kg) | 13<br>106<br>59<br>84<br>50                  | al is dit<br>surface<br>anners -   | •   |                   |  |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | mater<br>iety o<br>erent   |     |                   |  |
| Chemical<br>Name         | Sodium Dichromate                            | Disposition of this mater<br>It is used in a variety o<br>disposed of in different<br>to 1979. | · · |                   |  |

| Chemical Name: Phosphoric Acid   |
|--|
| üses:  |
| SolidLiquidXGas  |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Toxic by ingestion and inhalation. Irritant to skin and eyes. Tolerance - 1 mg per cubic meter of air. |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
| Level and Type of Worker Protection:   |
|  |
| Source of Chemical Usage Information:  |
| Source of Chemical Distribution Information/Accuracy:  |

## LSTINALLD CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |   |   |      |
|--------------------------|---|---|------|
| Other .                  |   |   |      |
| Offsite<br>Burial        |   |   |      |
| Onsite<br>Burial         |   |   |      |
| Onsite<br>Storage        |   |   |      |
| L. Iquid<br>Effluent     | ·   |   |      |
| Airborne<br>Release      |   | ed.<br>ery;   |      |
| Product                  |   | be estimated.<br>anium recovery<br>estimate<br>sing the H <sub>3</sub> <sup>20</sup><br>No  |      |
| Quantity<br>Used<br>(Kg) | 1,920<br>2,830<br>790<br>118<br>107<br>286<br>805 | ial canno<br>through u<br>but canno<br>rocedure<br>ely 1962.  | <br> |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980      | material e put thro rage, but Etch proce roximately prior to 19   |      |
| Shewical<br>Vame         | Phosphoric Acid                                   | Disposition of this material cannot be estimated. Some of it had to be put through uranium recovery; some to on-site storage, but cannot estimate relative amounts. Etch procedure using the H <sub>3</sub> PO was instituted approximately 1962. No records available prior to 1979. | · ·  |

| Chemical Name: Trichloroethylene  |
|---|
| Uses:   |
| SolidLiquid_XGas  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):                  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:                               |
| Toxic by inhalation, causes cancer in mice. Tolerance - 100 ppm in air. |
| Known or Suspected Environmental Impacts:                               |
|   |
|   |
|   |
| Level and Type of Worker Protection:                                    |
| •   |
| a a 1 Cara Information:   |
| Source of Chemical Usage Information:                                   |
| ·   |
| Source of Chemical Distribution Information/Accuracy:                   |
|   |

LSTINATED CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |   |
|--------------------------|---|
| 0ther                    |   |
| Offsite<br>Burial        |   |
| Onsite<br>Durial         |   |
| Ons I te<br>Storage      |   |
| l. Iquid<br>Effluent     | · · ·   |
| Airborne<br>Release      | -<br>20<br>20<br>20<br>20<br>20<br>20<br>20<br>20 |
| Product                  |   |
| Quantity<br>Used<br>(Kq) | 16<br>14<br>39<br>46<br>16<br>37                  |
| Year                     |   |
| Chemical                 | chloroethylene                                    |

| Chemical Name: Lime (Calcium Oxide)   |               |
|---|---------------|
| üses:   |               |
| Solid X Liquid Gas  | •             |
| Listed as Toxic: Yes No   |               |
| Status of Environmental Compliance (Past and Present):                              |               |
|   |               |
| Known or Suspected Public Health Impacts:   |               |
| Slightly dangerous on contact with water, steam, acid, or acid fumes. produce heat. | Will react to |
| Known or Suspected Environmental Impacts:   |               |
| Level and Type of Worker Protection:  |               |
| Source of Chemical Usage Information:   |               |
| Source of Chemical Distribution Information/Accuracy:                               |               |

LST HATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |                              |                                 |  |
|--------------------------|--|------------------------------|---------------------------------|--|
| Other .                  |  |                              |                                 |  |
| Offsite<br>Burial        |  |                              |                                 |  |
| Onsite<br>Burial         | ខេត្តបាន   | വവവ                          |                                 |  |
| Onsite<br>Storage        |  |                              | ·                               |  |
| L.Iquid<br>Effluent      | 95*<br>95<br>95<br>95<br>98<br>100<br>100                        | 95<br>95<br>95               | c um chloride                   |  |
| Airborne<br>Release      | ·  | •                            | e, and calc                     |  |
| Product                  |  |                              | calcium sulfate,                |  |
| Quantity<br>Used<br>(Kg) | 6,000<br>6,000<br>6,000<br>6,000<br>115,000<br>115,000<br>12,900 | 410,000<br>310,000<br>86,000 |                                 |  |
| Year                     | 1957<br>1956<br>1955<br>1954<br>1953<br>1952<br>1951             | 1946<br>1945<br>1944         | in mi                           |  |
| Shemical                 |  |                              | *Discharged as calcium nitrate, |  |

| Chemical Name: Hydrogen Peroxide   |
|--|
| üses:  |
| SolidLiquid_XGas   |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
| ·  |
| Known or Suspected Public Health Impacts:  |
| Concentrated solutions are highly toxic and strong irritants. Can easily cause blistering of the skin. Dangerous fire and explosion risk. Tolerance 1 ppm in air |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
|  |
| Level and Type of Worker Protection:   |
|  |
|  |
| Source of Chemical Usage Information:  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

### ESTINALLO CHEMICAL USAGE

### ALPHA AND BETA PROCESS

Estimated Distribution in %

| Chemical  | Year             | Quantity<br>Used<br>(Kg)               | Product                     | Airborne<br>Release                    | Liquid<br>Effluent | Onsite<br>Storage | Onsite<br>Burial | Offsite<br>Burial | 0ther | Unknown |
|---|------------------|--|-----------------------------|--|--------------------|-------------------|------------------|-------------------|-------|---------|
| gen Peroxide  |                  | 36,000<br>175,000<br>113,000<br>40,000 |                             | 10*<br>10<br>10                        | **06<br>06<br>06   |                   | ·                |                   |       |         |
| •   |                  |  |                             |  |                    |                   |                  |                   |       |         |
| *Converted to water, carbon monoxide, carbon by subsequent evaporation and pryrolysis of t precipitation process. | carbon<br>tion a | monoxide,<br>nd pryroly                | carbon diox<br>sis of the p | dioxide, oxygen, etc.,<br>the produced | an, etc.,          |                   |                  |                   |       |         |
| **Discharged as dilute nitric   | nitri            | c acid and                             | acid and hydrogen peroxide. | eroxide.                               |                    |                   |                  |                   |       |         |
| ٠,  |                  |  |                             |  |                    |                   |                  |                   |       |         |
|   |                  |  |                             |  |                    |                   |                  |                   |       |         |
|   |                  |  |                             |  |                    |                   |                  |                   |       |         |
|   |                  |  |                             |  |                    |                   |                  |                   |       |         |

| Chemical Name: Calcium Nitrate  |
|---|
| Uses:   |
| Solid X Liquid Gas .  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Strong oxidizer. Dangerous fire risk in contact with organic materials. May explode if shocked or heated. |
|   |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection:  |
|   |
| Source of Chemical Usage Information:   |
| Source of Chemical Usage Information.   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |
|   |

### LSTINALLD CHEMICAL USAGE

BETA PROCESS

Estimated Distribution in %

| Unknown                  |                       |   |   |      |
|--------------------------|-----------------------|---|---|------|
| Other                    |                       |   |   |      |
| Offsite<br>Burial        |                       |   |   |      |
| Ons ite<br>Burial        |                       |   |   |      |
| Ons I te<br>Storage      |                       | ·   |   |      |
| L. Iquid<br>Effluent     | 100*<br>100<br>100    |   |   |      |
| Airborne<br>Release      |                       | th nitric acid  |   |      |
| Product                  | ·                     | solution with   |   | <br> |
| Quantity<br>Used<br>(Kg) | 1,650<br>1,250<br>350 | 1y 40% sc   | • |      |
| Year                     | 1946<br>1945<br>1944  | oximate<br>inants   |   | <br> |
| Chemica 1<br>Name        | Calcium Nitrate       | *Discharged as approximately 40% iron nitrate contaminants. | • |      |

| Chemical Name: Diethyl Ether   |
|--|
| Uses:  |
| SolidLiquidXGas  |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
| •  |
| Known or Suspected Public Health Impacts:  |
| Extremely flammable. Severe fire and explosion hazard when exposed to heat or flame Forms explosive peroxides. Tolerance 400 ppm in air. |
| Known or Suspected Environmental Impacts:  |
|  |
| Level and Type of Worker Protection:   |
| Source of Chemical Usage Information:  |
| Source of Chemical Distribution Information/Accuracy:  |

## LSTINALLD CHEMICAL USAGE

BETA PROCESS

Estimated Distribution in 2

| Unknown                  |          |                           |  |                 |         |   |   |
|--------------------------|----------|---------------------------|--|-----------------|---------|---|---|
| <b>5</b>                 |          | <br>                      |  | - <del></del> - |         |   |   |
| Other .                  | *06      |                           |  |                 |         |   |   |
| Offsite<br>Burial        |          |                           |  |                 |         |   |   |
| Onsite<br>Burial         |          |                           |  | <br>            |         |   |   |
| Onsite<br>Storage        |          |                           |  |                 |         |   |   |
| Liquid<br>Effluent       |          |                           |  |                 |         |   |   |
| Airborne<br>Release      | 10       |                           |  |                 |         |   | · |
| Product                  |          |                           |  |                 |         | · |   |
| Quantity<br>Used<br>(Kg) | 3,000    | <br>yrolysis.             |  |                 | <u></u> |   |   |
| Year                     |          |                           |  |                 |         |   |   |
| Chemica 1<br>Name        | yl Ether | *Eventually decomposed by |  |                 |         |   |   |

| Chemical Name: Cupric Nitrate  |
|--|
| üses:  |
| SolidX Liquid Gas  |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Irritating to skin and mucous membrane. In contact with organic materials it may cause violent combustion and explosion. |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
| Level and Type of Worker Protection:   |
|  |
|  |
| Source of Chemical Usage Information:  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

## I STINALLD CHEMICAL USAGE

BETA PROCESS

Estimated Distribution in 2

| Unknown                  |                              |   |       |   |
|--------------------------|------------------------------|---|-------|---|
| Other .                  |                              |   |       |   |
| Offsite<br>Burial        |                              |   | ,<br> |   |
| Onsite<br>Aurial         |                              |   |       |   |
| Onsite<br>Storage        |                              |   |       |   |
| t. iquid<br>Effluent     | 100*<br>100                  |   |       |   |
| Airborne<br>Release      |                              | lution.   |       |   |
| Product                  |                              | n nitrate s   |       |   |
| Quantity<br>Used<br>(Kg) | 163,000<br>124,000<br>34,500 | ncentratio  |       |   |
| Year                     |                              | oo mn   |       | · |
| Chemica 1<br>Name        | ic Nitrate                   | *Discharged as a medium concentration nitrate solution. |       |   |

# LSTINALLD CHEMICAL USAGE

BETA PROCESS

Estimated Distribution in &

| Chemical<br>Name<br>Ethyl Alcohol         | - 1 | ty  | Product | rborne<br>lease | f. iquid<br>Effluent | Onsite<br>Storage | Ons ite<br>Burial | Offsite<br>Burial | Other<br>95* | Unknown |
|---|-----|-----|---------|-----------------|----------------------|-------------------|-------------------|-------------------|--------------|---------|
| 1945<br>1944<br>*Decomposed by pyrolysis. |     | 760 |         | റ ശ             | •                    | ·                 |                   |                   | 2<br>0<br>0  |         |
| •   |     |     |         |                 |                      |                   |                   |                   |              | ,       |
|   |     |     |         | <u> </u>        |                      |                   |                   |                   |              |         |

| Chemical Name: Carbon Tetrachloride  |
|--|
| Uses:  |
| Solid Liquid X Gas   |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
| Known or Suspected Public Health Impacts: It can be toxic by inhalation, injestion or skin absorption. Acute effects are nausea, vomiting, diarrhea, headache, stupor, renal and liver damage, and possibly death. Chronic effects are primarily liver damage but kidney inury and visual disturbances also occur. Skin contact can lead to dermatitis through defatting action. According to IARC, evidence for carcinogenicity to humans is inadequate; evidence is sufficient for animals.  Known or Suspected Environmental Impacts: |
| Level and Type of Worker Protection:  Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respirate protection.  Source of Chemical Usage Information:  |
| Source of Chemical Distribution Information/Accuracy:  |

L'STIMATED CHEMICAL USAGE

ALPHA AND BETA PROCESS

Estimated Distribution in X

| Unknown                  |   |   |   |      |  |
|--------------------------|---|---|---|------|--|
| Other                    |   |   |   |      |  |
| Offsite<br>Burial        |   |   |   | ,    |  |
| Onsite<br>Burial         |   |   |   | <br> |  |
| Ons (te<br>Storage       |   |   |   |      |  |
| l. Iquid<br>Effluent     | 99<br>99<br>99                          |   |   |      |  |
| Airborne<br>Release      |   |   | carbonate                                 |      |  |
| Product                  | 33<br>33<br>33<br>33                    |   | sodium Nydroxide to                       |      |  |
| Quantity<br>Used<br>(Kg) | 57,300<br>668,500<br>717,500<br>261,800 | - |   |      |  |
| Year                     | 1946<br>1945<br>1944<br>1943            |   | onia or                                   |      |  |
| Chemica l<br>Name        | Carbon Tetrachloride                    |   | *Neutralized with ammonia<br>or chloride. |      |  |

| Chemical Name: Phosgene   |
|---|
| Uses:   |
| Solid Liquid X Gas X  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Highly toxic by inhalation, strong irritant to eyes, tolerance - 0.1 ppm in air. May cause severe pulmonary edema or pneumonia. Inhalation of high concentrations causes choking, constricted feeling in chest, coughing, painful breathing.  Known or Suspected Environmental Impacts: |
| Level and Type of Worker Protection:  |
| Supplied air, full face mask.   |
|   |
| Source of Chemical Usage Information:   |
| Source of Charical Distribution Information/Accuracy:   |

LSTIMALLD CHEMICAL USAGE

ALPHA AND BETA PROCESS

Estimated Distribution in 2

| Chemical<br>Name      | Year       | Quantity<br>Used<br>(Kg)      | Product                                     | Airborne<br>Release            | L. Iquid<br>Effluent | Onsite<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other | Unknown |
|-----------------------|------------|-------------------------------|---|--------------------------------|----------------------|-------------------|------------------|-------------------|-------|---------|
| Phosgene              |            | 201,500<br>475,500<br>175,000 |   | 5 5 5<br>5 6 7<br>5 7<br>7 8 7 | **<br>86<br>86       |                   |                  |                   |       |         |
| *Neutralized with am  | onia t     | o carbonat                    | e and chlor                                 | de.                            |                      |                   |                  |                   |       |         |
| **Neutralized with so | ium hy     | droxide to                    | sodium hydroxide to carbonate and chloride. | and chlorid                    | ů                    |                   |                  |                   |       |         |
|                       | <u>.</u> . |                               |   |                                |                      | ·                 |                  |                   |       |         |
|                       |            |                               |   |                                |                      |                   |                  |                   |       |         |
|                       |            |                               |   |                                |                      |                   |                  |                   |       |         |
|                       |            |                               |   |                                |                      |                   |                  |                   |       |         |
|                       |            |                               | •   |                                |                      |                   |                  |                   |       |         |

| Chemical Name: Ammonia   |
|--|
| Uses:  |
| SolidLiquidXGas  |
| Listed as Toxic: YesNo   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
| •  |
| Known or Suspected Public Health Impacts:  |
| Toxic and irritant by inhalation. Tolerance 25 ppm in air. Inhalation of concentrate fumes may be fatal. moderate fire risk. |
| Known or Suspected Environmental Impacts:  |
|  |
| Level and Type of Worker Protection:   |
| Source of Chemical Usage Information:  |
| Source of Chemical Distribution Information/Accuracy:  |

LSLINALLD CHEMICAL USAGE

ALPHA AND BETA PROCESS

Estimated Distribution in &

| Unknown                  |                                     |   |   |      |
|--------------------------|-------------------------------------|---|---|------|
| Other                    |                                     |   |   |      |
| Offsite<br>Burial        |                                     |   |   |      |
| Onsite<br>Aurial         |                                     |   |   |      |
| Onsite<br>Storage        |                                     |   | · |      |
| L. Iquid<br>Effluent     | 955 <del>*</del>                    | ate,  |   |      |
| Airborne<br>Release      | വവവ                                 | , sodium carbonate,   |   |      |
| Product                  |                                     | hloride, so<br>um nitrate.  |   |      |
| Quantity<br>Used<br>(Kg) | 8,500<br>61,500<br>42,000<br>15,000 | f sodium c<br>ute ammon'  |   |      |
| Year                     |                                     | tion c  |   | <br> |
| Chemica 1<br>Name        | ia                                  | *Discharged as a solution of sodium dhloride, sodium hypochlorite, or dilute ammonium nitra | · |      |

| Chemical Name: Sulfuric Acid  |
|---|
| Uses:   |
| SolidLiquid_XGas  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:  It is corrosive to all body tissues. Inhalation of concentrated vapor may cause seridamage to lungs. Contact with eyes may result in total loss of vision; skin contact may produce severe necrosis. Frequent contact with dilute solution can cause dermatitis. Ingestion may cause severe injury and death.  Known or Suspected Environmental Impacts:  Presence of atmospheric sulfuric acid can contribute to the acidification of rain water which affects animal and plant life. |
| Level and Type of Worker Protection: Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation (2) rubber clothing (3) chemical goggles (4) face shield (5) proper respiratory protection.  |
| Source of Chemical Usage Information:   |
|   |
|   |

Source of Chemical Distribution Information/Accuracy:

# L'STINATED CHEMICAL USAGE

Estimated Distribution in 2

| Unknown                  |                        |  |   |   |       |  |
|--------------------------|------------------------|--|---|---|-------|--|
| Other .                  |                        |  |   |   |       |  |
| Offsite<br>Burial        |                        |  |   |   |       |  |
| Onsite<br>Burial         | 50<br>50               |  |   |   |       |  |
| Onsite<br>Storage        |                        |  |   |   |       |  |
| L. Iquid<br>Effluent     | 50*<br>50              |  |   |   |       |  |
| Airborne<br>Release      |                        | calcium sulfate                                |   |   |       |  |
| Product                  |                        |  |   |   |       |  |
| Quantity<br>Used<br>(Kg) | 1,500,000<br>1,500,000 | acid concentration of                          |   |   | ····· |  |
| Year                     |                        |  |   |   |       |  |
| Chemica 1<br>Name        | Sulfuric Acid          | *Discharged as a weak<br>or magnesium sulfate. | • | • |       |  |

| Chemical Name: Manganese Dioxide                             |
|--|
| Uses:  |
| Solid X Liquid Gas .   |
| Listed as Toxic: Yes No                                      |
| Status of Environmental Compliance (Past and Present):       |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:                    |
| Oxidizing agent. May ignite organic materials. Mildly toxic. |
| ·  |
| Known or Suspected Environmental Impacts:                    |
|  |
|  |
|  |
| Level and Type of Worker Protection:                         |
|  |
|  |
| Source of Chemical Usage Information:                        |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:        |

LSTINALLD CHENICAL USAGI.

Estimated Distribution in &

| Unknown                  |                      | , and the second |                               |                                       |      |              |
|--------------------------|----------------------|------------------|-------------------------------|---------------------------------------|------|--------------|
| Other                    |                      |                  |                               |                                       |      |              |
| Offsite<br>Burial        |                      |                  |                               |                                       |      |              |
| Ons ite<br>Burial        | 95*<br>95            |                  |                               |                                       | <br> |              |
| Onsite<br>Storage        |                      |                  |                               |                                       |      |              |
| L. Iquid<br>Effluent     | വ                    |                  |                               |                                       |      |              |
| Airborne<br>Release      |                      |                  |                               |                                       |      | <del></del>  |
| Product                  |                      |                  |                               | · · · · · · · · · · · · · · · · · · · |      |              |
| Quantity<br>Used<br>(Kg) | 24,000<br>24,000     |                  | o                             |                                       |      | <br>         |
| Year                     | 1959<br>1958<br>1958 | •                | sulfat<br>                    |                                       |      | <br><u>_</u> |
| Chemica 1<br>Name        | Manganese Dioxide    |                  | *Buried as manganese sulfate. |                                       |      |              |

| Chemical Name: Ammonium Sulfate                        |
|--|
| Uses:  |
| SolidX Liquid Gas                                      |
| Listed as Toxic: YesNo                                 |
| Status of Environmental Compliance (Past and Present): |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:              |
| Low toxicity, mild irritant.                           |
|  |
| Known or Suspected Environmental Impacts:              |
| ·  |
|  |
|  |
| Level and Type of Worker Protection:                   |
|  |
| ·  |
| Source of Chemical Usage Information:                  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |

L'STINATED CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |                        |                                |      |      |  |
|--------------------------|------------------------|--------------------------------|------|------|--|
| Other                    |                        |                                |      |      |  |
| Offsite<br>Burial        |                        |                                |      |      |  |
| Onsite<br>Burial         |                        |                                | <br> | <br> |  |
| Onsite<br>Storage        |                        |                                | <br> |      |  |
| l. Iquid<br>Effluent     | 100*<br>100            |                                |      |      |  |
| Airborne<br>Release      |                        |                                |      |      |  |
| Product                  |                        | olution.                       |      |      |  |
| Quantity<br>Used<br>(Kg) | 1,400,000<br>1,400,000 | a weak sulfuric acid solution. |      |      |  |
| Year                     |                        | sulfu                          |      | <br> |  |
| Chemical<br>Name         | Ammonium Sulfate       | *Discharged as                 |      |      |  |

| Chemical Name: Belguard 550  |
|--|
| Uses:  |
| SolidLiquid_X Gas  |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| May cause skin and eye irritation.   |
|  |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
|  |
| Level and Type of Worker Protection:   |
| Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include |
| (1) adequate ventilation, (2) rubber gloves, (3) chemical goggles (4) proper respirat protection.  |
| Source of Chemical Usage Information:  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

ESTIMATED CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  | ·                                | <u>.</u> |       |  | <br> |   | ·    |             |
|--------------------------|----------------------------------|----------|-------|--|------|---|------|-------------|
| Other                    |                                  |          |       |  |      |   |      |             |
| Offsite<br>Burial        |                                  |          |       |  |      |   |      |             |
| Onsite<br>Burial         |                                  |          |       |  | <br> |   |      |             |
| Onsite<br>Storage        |                                  |          |       |  |      |   |      |             |
| Liquid<br>Effluent       | 100%                             | 100%     | 100%  |  |      |   |      |             |
| Airborne<br>Release      |                                  |          |       |  |      |   |      | <del></del> |
| Product                  | ·                                |          |       |  |      |   |      |             |
| Quantity<br>Used<br>(Kg) | 000°6                            | 55,000   | 5,000 |  |      | - | <br> |             |
| Year                     |                                  | 1985     | 1984  |  |      |   | <br> |             |
| Chemical<br>Name         | Belguard 550<br>(Bell Chem. Co.) |          |       |  |      |   |      |             |

| Chemical Name: Biotreat VIII  |
|---|
| Uses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| It is a severe eye irritant and causes dry skin to redden and burn.   |
|   |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include |
| (1) adequate ventilation (2) rubber gloves, (3) chemical goggles (4) proper respirato protection.   |
| Source of Chemical Usage Information:   |
|   |
| Source of Chamical Dispublican Todayananian (laguage)   |
| Source of Chemical Distribution Information/Accuracy:   |

LSTINATED CHEMICAL HSAGE

Estimated Distribution in X

| Year   | Quantity<br>Used<br>(Kg) | Product    | Airborne<br>Release | Liquid<br>Effluent | Onsite<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other . | · Unknown |
|--|--------------------------|------------|---------------------|--------------------|-------------------|------------------|-------------------|---------|-----------|
| Biotreat VIII 1986<br>(Bell Chem. Co.) (thru | 5 1,000<br>ru<br>n)      |            |                     | 100%               |                   |                  |                   |         |           |
| 1985   | 2 6,000                  |            |                     | 100%               |                   |                  |                   |         |           |
| 1984   | 4 550                    |            |                     | 100%               |                   |                  |                   |         |           |
|  |                          |            |                     |                    |                   |                  |                   |         |           |
|  |                          |            |                     |                    |                   |                  |                   |         |           |
|  |                          |            |                     |                    |                   |                  |                   |         |           |
|  |                          |            |                     |                    |                   |                  |                   |         |           |
| <del>- ; -</del>                             | -                        | · <u>·</u> |                     |                    |                   |                  |                   |         |           |
|  |                          |            |                     |                    |                   |                  |                   |         |           |
|  |                          |            | <del></del>         | _                  |                   |                  |                   |         |           |
|  | <del></del>              |            |                     |                    |                   |                  |                   |         | ·         |
|  |                          |            | <del></del>         |                    |                   |                  |                   |         |           |

| Chemical Name: Wrico ZC-304   |
|---|
| Uses:   |
| SolidLiquidGas  |
| Listed as Toxic: Yes No   |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
|   |
| ·   |
| The Common of Engineers I Imports:  |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
| ·   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) chemical goggles, (4) proper respirate protection. |
| Source of Chemical Usage Information:   |
| COULCE OF CHEMICAL COAGE INTOLMACTON.   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |
|   |

| Chemical Name: Zimmite ZC-304   |
|---|
| Uses: Corrosion inhibitor   |
| Solid Liquid X Gas  |
| Listed as Toxic: Yes No   |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| It is mildly toxic when ingested and eye and skin irritant.   |
|   |
| Known or Suspected Environmental Impacts:   |
| •   |
|   |
|   |
| Level and Type of Worker Protection:  |
| Specific use, conditions, quantities and other factors determine the level and type   |
| of worker protection. Standard measures for protection against this material include (1) adequate ventilation (2) rubber gloves, (3) chemical goggles, (4) proper respirate |
| protection.   |
| Source of Chemical Usage Information:   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |
|   |

## ISTINATED CHEMICAL USAGE

Estimated Distribution in 2

| Chemica 1<br>Name                                | Year | Quantity<br>Used<br>(Kg) | Product | Airborne<br>Release | Liquid<br>Effluent | Ons I te<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other . | Unknown         |
|--|------|--------------------------|---------|---------------------|--------------------|---------------------|------------------|-------------------|---------|-----------------|
| Wrico ZC-304*<br>(Wright Chem. Co.)              | 1985 | 006                      |         |                     | 100%               |                     |                  |                   |         |                 |
| Zimmite ZC-304*<br>(Zimmite Chem. Co.)           | 1984 | 55,000                   |         |                     | 100%               | ,                   |                  |                   |         |                 |
|  | 1983 | 55,000                   |         |                     | 100%               |                     |                  |                   |         | -               |
| (1980-Feb. 1985 used 1982 in all cooling towers) |      | 55,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
| (1974-1980 used in<br>one half of towers)        | 1981 | 55,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
|  | 1980 | 55,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
|  | 1979 | 28,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
|  | 1978 | 28,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
| *Note: Wrico 2C-304 1977                         |      | 28,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
| and finalite 20-304<br>are the same              | 1976 | 28,000                   |         | _                   | 100%               |                     |                  |                   |         |                 |
| Chemical; wright<br>Chemical Company             | 1975 | 28,000                   |         |                     | 100%               |                     |                  |                   |         |                 |
| Dougnt the Limmite<br>Chemical Company.          | 1974 | 28,000                   |         |                     | 100%               |                     |                  |                   |         | -: <del>-</del> |
|  |      |                          |         | <del></del>         |                    |                     |                  |                   |         | ·               |

| Chemical Name: Chemtrol 19   |
|--|
| Uses: Microbiocide, for cool tower water treatment   |
| SolidLiquidX Gas   |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
| <del>.</del>   |
| Known or Suspected Public Health Impacts:  |
| It can be harmful if swallowed and is a mild eye and skin irritant.  |
|  |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
| ·  |
| Level and Type of Worker Protection:   |
| Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) chemical goggles, (4) proper respirate protection. |
| Source of Chemical Usage Information:  |
|  |
|  |

Source of Chemical Distribution Information/Accuracy:

## L'STINATED CHEMICAL USAGE

Estimated Distribution in %

| . Unknown                |                 |   | <u> </u> |  |        |        |                    |                      |        |        |        | -, -,  |          |  |
|--------------------------|-----------------|---|----------|--|--------|--------|--------------------|----------------------|--------|--------|--------|--------|----------|--|
| 0ther                    |                 |   |          |  |        |        |                    |                      |        |        |        |        |          |  |
| Offsite<br>Burial        |                 |   |          |  |        |        |                    |                      |        |        |        | •      |          |  |
| Onsite<br>Burial         |                 |   |          |  |        |        |                    |                      |        |        |        |        |          |  |
| Onsite<br>Storage        |                 | ····                                      |          |  |        |        |                    |                      |        |        |        |        | <u> </u> |  |
| l.Iquid<br>Effluent      | 100%            | 100%                                      | 100%     | 100%   | 100%   | 100%   | 100%               | 100%                 | 100%   | 100%   | 100%   | 100%   | <br>     |  |
| Airborne<br>Release      |                 |   |          |  |        |        |                    |                      |        |        |        |        |          |  |
| Product                  |                 |   |          |  |        |        |                    |                      |        |        |        |        |          |  |
| Quantity<br>Used<br>(Kg) | 400             | 22,000                                    | 22,000   | 22,000   | 22,000 | 22,000 | 11,000             | 11,000               | 11,000 | 11,000 | 11,000 | 11,000 |          |  |
| Year                     | 1985            | 1984                                      | 1983     |  | 1981   | 1980   | 1979               | 1978                 | 1977   | 1976   | 1975   | 1974   |          |  |
| Chemica]<br>Name         | Chemtrol No. 19 | (Wright or Zimmite<br>Chemical Companies) |          | (1980-Feb. 1985 used 1982 in all cooling towers) |        |        | (1974-1980 used in | Olle Hall Of COWERS) |        |        |        |        |          |  |

| Chemical Name: Endcor 4622  |     |
|---|-----|
| Uses: Cooling Water Treatment   |     |
| SolidLiquidX Gas  |     |
| Listed as Toxic: YesNo  |     |
| Status of Environmental Compliance (Past and Present):  |     |
|   |     |
|   |     |
|   |     |
| Known or Suspected Public Health Impacts:   |     |
| It will irritate eyes on contact and may irritate skin on prolonged contact   | ot. |
|   |     |
| Known or Suspected Environmental Impacts:   |     |
|   |     |
|   |     |
|   |     |
| Level and Type of Worker Protection:  |     |
| Specific use, conditions, quantities, and other factors determine the level of worker protection. Standard measures for protection against this mater |     |
| (1) adequate ventilation (2) rubber gloves, (3) chemical goggles, (4) proprespiratory protection.   |     |
| Source of Chemical Usage Information:   |     |
|   |     |
|   |     |
| Source of Chemical Distribution Information/Accuracy:   |     |

# L'STIMATED CHEMICAL USAGE

Estimated Distribution in &

| Chemical<br>Name       | Year | Quantity<br>Used<br>(Kg) | Product | Airborne<br>Release | l.iquid<br>Effluent | Onsite<br>Storage | Onsite<br>Burial | Offsite<br>Burial | Other      | Unknown |
|------------------------|------|--------------------------|---------|---------------------|---------------------|-------------------|------------------|-------------------|------------|---------|
| Encor 4622             | 1980 | 28,000                   |         |                     | 100%                |                   |                  |                   |            |         |
| (Dearborn Cheill: CO.) | 1979 | 28,000                   |         |                     | 100%                |                   |                  |                   | . <u> </u> |         |
|                        | 1978 | 28,000                   |         |                     | . %001              |                   |                  |                   |            |         |
|                        | 1977 | 28,000                   |         |                     | 100%                |                   |                  |                   |            |         |
|                        | 1976 | 28,000                   |         |                     | 100%                |                   |                  |                   |            |         |
|                        | 1975 | 28,000                   |         |                     | 100%                |                   |                  |                   |            |         |
| ·                      | 1974 | 28,000                   |         |                     | 100%                |                   |                  |                   |            |         |
|                        |      |                          |         |                     |                     |                   | •                |                   |            |         |
|                        |      |                          |         |                     |                     |                   |                  |                   |            |         |
|                        |      |                          |         |                     |                     |                   |                  | -                 |            |         |
|                        |      | :                        |         |                     |                     | <u></u>           |                  |                   |            |         |
|                        |      |                          |         | . <del></del>       |                     |                   |                  |                   |            |         |
|                        |      |                          |         |                     |                     |                   |                  |                   |            |         |
|                        |      | ·                        |         |                     |                     |                   |                  |                   |            |         |
|                        |      |                          |         | <del></del>         |                     |                   |                  |                   |            |         |

| Chemical Name: Microbiotreat 321  |
|---|
| Uses: Biocide   |
| SolidLiquidXGas   |
| Listed as Toxic: Yes No   |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Eye and skin irritant.  |
|   |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection:<br>Specific use, conditions, quantities, and other factors determine the level and type  |
| of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) chemical goggles, (4) proper |
| respiratory protection.   |
| Source of Chemical Usage Information:   |
|   |
|   |

Source of Chemical Distribution Information/Accuracy:

# LSTINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                   |        |        |        |          |        |        |       |      |   |   |      |                 |
|--------------------------|-------------------|--------|--------|--------|----------|--------|--------|-------|------|---|---|------|-----------------|
| Other .                  |                   |        |        |        |          |        |        |       |      |   |   |      |                 |
| Offsite<br>Burial        |                   |        |        |        |          |        |        |       |      |   |   |      | _               |
| Onsite<br>Burial         |                   |        |        |        |          |        |        |       |      |   |   | <br> | <br>_           |
| Onsite<br>Storage        |                   |        |        |        |          |        |        |       | <br> |   |   |      | <br>            |
| Llquid<br>Effluent       | 100%              | 100%   | 100%   | 100%   | 100%     | 100%   | 100%   |       |      |   |   | <br> |                 |
| Airborne<br>Release      |                   |        |        |        |          |        |        |       |      |   |   | <br> | <br><del></del> |
| Product                  |                   |        |        |        |          |        |        |       |      |   |   | <br> |                 |
| Quantity<br>Used<br>(Kg) | 11,000            | 11,000 | 11,000 | 11,000 | 11,000   | 11,000 | 11,000 |       |      | - | _ |      |                 |
| Year                     | 1980              | 1979   | 1978   | 1977   | 1976     | 1975   | 1974   | <br>- |      |   | • |      | <br>            |
| Chemica 1                | Microbiotreat 321 |        |        |        | TARLES A |        |        |       | -    |   |   |      | _               |

| Chemical Name: HTH   |
|--|
| Uses:  |
| Solid Gas  |
| Listed as Toxic: YesNo   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
|  |
|  |
| Known or Suspected Environmental Impacts:  |
|  |
|  |
|  |
| To the difference of Western Dropper Inc.  |
| Level and Type of Worker Protection:   |
| Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection. |
| Source of Chemical Usage Information:  |
|  |
| To Some tion (local page)  |
| Source of Chemical Distribution Information/Accuracy:  |

# LSTIMALLD CHIMICAL USAGE

Estimated Distribution in &

|               |      | Quantity |         |         |          | 246000  |         | 0 4 4 5 9 9 0 |       |         |
|---------------|------|----------|---------|---------|----------|---------|---------|---------------|-------|---------|
| Name          | Year | (Kg)     | Product | Release | Effluent | Storage | Our ial | Burial        | Other | Unknown |
| HTH (Calcium  | 1973 | 300      |         |         | 100%     |         |         |               |       |         |
| hypochiorite) | 1972 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1971 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1970 | 300      |         |         | 100%     |         |         | <del></del>   |       |         |
|               | 1969 | 300      | -       |         | 100%     |         |         |               |       |         |
|               | 1968 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1967 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1966 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1965 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1964 | 300      |         |         | 100%     |         |         |               |       |         |
|               | 1963 | 300      |         |         | 100%     |         |         |               |       |         |
|               |      |          |         |         |          |         |         |               |       |         |
|               |      |          |         |         |          |         |         |               |       |         |
|               |      |          |         |         |          |         |         |               |       |         |
|               |      |          | •       |         |          |         |         |               |       |         |
|               |      |          |         |         |          |         |         |               |       |         |

| Chemical Name: Freons  |
|--|
| Uses:  |
| SolidLiquidX GasX  |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| High vapor concentrations may cause confusion, pulmonary irritation, tremors and rarely coma. These effects are transient. |
| Tarety Onk. These effects are transfer.  |
| Known or Suspected Environmental Impacts:  |
| •  |
|  |
|  |
| Level and Type of Worker Protection:   |
| Level and Type of world from   |
|  |
| Source of Chemical Usage Information:  |
| Source of Chemical Usage Information.  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

# L'STIMATED CHEMICAL USAGE

Estimated Distribution in %

| Chemica)  |               | Quantity<br>Used       |         | Airborne    | l. Iquid    | Ons i te |        | 0ffsite |       |         |
|---|---------------|------------------------|---------|-------------|-------------|----------|--------|---------|-------|---------|
| Name  | Year          | (Kg)                   | Product | [ ]         | Effluent    | Storage  | Durial | Burial  | 0ther | Unknown |
| Freons  | 1943-<br>1981 | ;<br>17,000<br>Kg/year |         |             |             |          |        |         | 100%* |         |
|   | 1981-<br>1985 | 17,000<br>Kg/year      |         |             |             | 100%     |        |         |       |         |
|   | 1986          | 4,000                  |         |             |             | 100%     |        |         |       |         |
|   |               |                        |         |             |             |          |        |         |       |         |
|   |               |                        |         |             |             |          |        |         |       |         |
| *Poured on ground.<br>Unknown quantity<br>excaped to the air. |               |                        |         |             |             |          |        |         |       |         |
|   |               |                        |         |             |             |          |        |         |       |         |
| 255   |               |                        |         |             |             |          |        |         |       |         |
|   |               |                        |         |             |             |          |        |         |       |         |
| Nove Frank  |               |                        |         |             |             |          |        |         |       |         |
|   |               |                        |         |             | <del></del> |          |        |         |       |         |
|   |               |                        |         | <del></del> |             |          |        |         |       |         |

| Chemical Name: Sulfur Dioxide  |
|--|
| Uses:  |
| Solid X Liquid Gas   |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Toxic by inhalation. Strong irritant to eyes and mucous membranes. Dangerous air contaminant and constituent of smog.  |
|  |
| Known or Suspected Environmental Impacts:  |
| ·  |
|  |
|  |
| Level and Type of Worker Protection:   |
|  |
|  |
| Source of Chemical Usage Information:  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
| Source of Chemical Distribution into the same and the sam |

## LSTINALLD CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |                 |             |
|--------------------------|-----------------|-------------|
| 0ther                    |                 |             |
| Offsite<br>Burial        |                 |             |
| Onsite<br>Burial         |                 |             |
| Onsite<br>Storage        |                 |             |
| Liquid<br>Effluent       |                 |             |
| Airborne<br>Release      | 100%            | <del></del> |
| Product                  |                 |             |
| Quantity<br>Used<br>(Kg) | 1955- 4,790,000 |             |
| Year                     | 1955-<br>1986   |             |
| Chemical                 | Sulfur Dioxide  |             |

| Chemical Name: Indene                                  |
|--|
| Uses:  |
| SolidLiquidX Gas                                       |
|  |
| Listed as Toxic: Yes                                   |
| Status of Environmental Compliance (Past and Present): |
|  |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:              |
| Low toxicity   |
|  |
| Known or Suspected Environmental Impacts:              |
| Known or Suspected Envilonmental impasts.              |
|  |
|  |
|  |
| Level and Type of Worker Protection:                   |
| Level and Type Of Mornor Troposition                   |
|  |
| ·  |
| Source of Chemical Usage Information:                  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |

## ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |  |      |  |
|--------------------------|--|--|------|--|
| Other                    |  |  |      |  |
| Offsite<br>Burial        |  |  |      |  |
| Onsite<br>Burial         |  |  |      |  |
| Onsite<br>Storage        | 40<br>55<br>80<br>80<br>80   |  | <br> |  |
| Liquid<br>Effluent       |  |  |      |  |
| Airborne<br>Release      |  |  |      |  |
| Product                  | 60<br>45<br>20<br>20<br>20   |  | <br> |  |
| Quantity<br>Used<br>(Kg) | 90<br>551<br>798<br>483<br>183                                     |  |      |  |
| Year                     | 1972<br>1971<br>1970<br>1969<br>1968                               |  |      |  |
| Chemical<br>Name         | Indene (in<br>preparation of<br>pitch and/or<br>carbon precursors) |  |      |  |

| Chemical Name: Cinnamic Aldehyde   |
|--|
| üses:  |
| SolidLiquidX Gas   |
| Listed as Toxic: Yes No  |
| Status of Environmental Compliance (Past and Present):   |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:  |
| Reactive compound, participating in oxidation, reduction, addition and polymerization reactions. |
| Known or Suspected Environmental Impacts:  |
| Level and Type of Worker Protection:   |
| Source of Chemical Usage Information:  |
| Source of Chemical Distribution Information/Accuracy:  |

ESTINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |
|--------------------------|--|
| 0ther                    |  |
| Offsite<br>Burial        |  |
| Onsite<br>Burial         | •  |
| Onsite<br>Storage        | 40   |
| l.iquid<br>Effluent      |  |
| Airborne<br>Release      |  |
| Product                  | 09   |
| Quantity<br>Used<br>(Kg) | 238  |
| Year                     |  |
| Chemical<br>Name         | Cinnamic aldehyde (pitch and carbon precursor) |

| Chemical Name: Ethylene dibromide   |
|---|
| Uses:   |
| SolidLiquidXGas   |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Toxic by inhalation, ingestion, or skin absorption. May be carcinogenic. Strong irritant to eyes and skin. Tolerence 20 ppm in air. |
| Known or Suspected Environmental Impacts:   |
|   |
| Level and Type of Worker Protection:  |
|   |
|   |
| Source of Chemical Usage Information:   |
|   |
| Source of Chemical Distribution Information/Accuracy:   |

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                    |
|--------------------------|--------------------|
| Other                    |                    |
| Offsite<br>Burial        |                    |
| Onsite<br>Burial         | 66                 |
| Onsite<br>Storage        |                    |
| Liquid<br>Effluent       |                    |
| Airborne<br>Release      |                    |
| Product                  |                    |
| Quantity<br>Used<br>(Kg) | 11,300             |
| Year                     |                    |
| Chemical<br>Name         | Ethylene Dibromide |

| Chemical Name: Urethane Resins                                |
|---|
| Uses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):        |
|   |
| Known or Suspected Public Health Impacts: Toxic by ingestion. |
| Known or Suspected Environmental Impacts:                     |
| Level and Type of Worker Protection:                          |
| Source of Chemical Usage Information:                         |
| Source of Chemical Distribution Information/Accuracy:         |

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| r Unknown                |   |  |  |
|--------------------------|---|--|--|
| Other .                  |   |  |  |
| Offsite<br>Burial        | •   |  |  |
| Onsite<br>Burial         |   |  |  |
| Onsite<br>Storage        |   | drds.  |  |
| Liquid<br>Effluent       |   | but no records.  |  |
| Airborne<br>Release      | 0.222222  | prior to 1979  |  |
| Product                  | 96<br>96<br>96<br>96  | naterial   |  |
| Quantity<br>Used<br>(Kg) | 5,727<br>4,295<br>5,523<br>5,114<br>4,295<br>4,500<br>5,523 | of this<br>on-site.                                      |  |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1980<br>1979        | le use<br>buriec   |  |
| Chemical<br>Name         | hane Foam<br>onent<br>olyol)                                | There was considerable use<br>Considerable amount buriec |  |

| Chemical Name: Ammonium Carbonate                      |
|--|
| Uses:  |
| Solid X Liquid Gas                                     |
| Listed as Toxic: Yes                                   |
| Status of Environmental Compliance (Past and Present): |
|  |
|  |
| •  |
| Known or Suspected Public Health Impacts:              |
| Respiratory irritant, source or ammonia.               |
|  |
| Known or Suspected Environmental Impacts:              |
|  |
|  |
|  |
| Level and Type of Worker Protection:                   |
|  |
|  |
| Source of Chemical Usage Information:                  |
|  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |

LSTINALLD CHEMICAL USAGE

ALPHA PROCESS

Estimated Distribution in %

| Unknown                  |                  |                      |      |   | <br> | · |
|--------------------------|------------------|----------------------|------|---|------|---|
| Other .                  |                  |                      |      |   | <br> |   |
| Offsite<br>Burial        |                  |                      |      |   |      |   |
| Onsite<br>Durial         |                  |                      | <br> |   |      |   |
| Onsite<br>Storage        |                  |                      |      |   | <br> |   |
| L. Iquid<br>Effluent     | 100*<br>100      |                      |      |   | <br> |   |
| Airborne<br>Release      |                  |                      |      |   | <br> |   |
| Product                  |                  |                      |      |   | <br> |   |
| Quantity<br>Used<br>(Kg) | 45,000<br>20,000 | trate.               |      |   |      |   |
| Year                     |                  | ammorium nitrate.    |      |   |      |   |
| Chemica)<br>Name         | Carbonate        | *Discharged as ammor |      | • | ·    |   |

| Chemical | Name:    | 1,1,1 - 5 | Prichloroethane | <b>e</b>      |     |           |  |
|----------|----------|-----------|-----------------|---------------|-----|-----------|--|
| Uses:    |          |           |                 |               |     |           |  |
| Solid    | Liqu     | idX       | Gas             | <del></del> · |     |           |  |
| Listed a | s Toxic: | Yes       | . No            |               |     |           |  |
| Status o | f Enviro | nmental   | Compliance      | (Past         | and | Present): |  |

Known or Suspected Public Health Impacts:

It is irritating to the eyes, mucous membranes, and in high concentrations, CNS depressant. Prolonged or repeated contact with skin results in transient erythema and defatting of skin. Mild liver and kidney dysfunction may occur transiently following recovery from CNS depression.

Known or Suspected Environmental Impacts:

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respiratory protection.

Source of Chemical Usage Information:

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  |   |
|--------------------------|--|---|
| Other                    |  | _ |
| Offsite<br>Burial        |  |   |
| Onsite<br>Burial         |  | _ |
| Onsite<br>Storage        | 4 m m m m  |   |
| l.iquid<br>Effluent      |  |   |
| Airborne<br>Release      | 97<br>97<br>97<br>97                                     |   |
| Product                  |  |   |
| Quantity<br>Used<br>(Kg) | 71,997<br>85,467<br>74,484<br>56,836<br>59,182<br>50,898 |   |
| Year                     | 1985<br>1984<br>1983<br>1982<br>1981<br>1979             |   |
| Chemical<br>Name         | 1,1,1 Tri-<br>chloroethane                               |   |

# ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                                       |
|--------------------------|---------------------------------------|
| Other                    |                                       |
| Offsite<br>Burial        |                                       |
| Onsite<br>Burial         |                                       |
| Onsite<br>Storage        | 1000%<br>1000%<br>1000%<br>1000%      |
| l.iquid<br>Effluent      |                                       |
| Airborne<br>Release      | -                                     |
| Product                  |                                       |
| Quantity<br>Used<br>(Kq) | 0<br>1.5<br>1.9<br>1.5<br>20.1        |
| Year                     |                                       |
| Chemical<br>Name         | chlorinated<br>enyls in<br>ng coolant |

L'STIMATED CHEMICAL USAGE

Estimated Distribution in %

| Other Unknown            |                                       |  |
|--------------------------|---------------------------------------|--|
| Offsite<br>Burial Ot     | ·                                     |  |
| Onsite<br>Burial         |                                       |  |
| Onsite<br>Storage        |                                       |  |
| Liquid .<br>Effluent     |                                       |  |
| Airborne<br>Release      | 0000                                  |  |
| Product                  | 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 |  |
| Quantity<br>Used<br>(Kg) | 307<br>307<br>307<br>409              |  |
| Year                     | 1983<br>1982<br>1979<br>1974<br>1971  |  |
| Chemical<br>Name         | M-Phenylene Diamine (MPDA)            |  |

| Chemical Name: M. D. A. (Methylenedianiline)           |
|--|
| Uses:  |
| Solid X Liquid Gas                                     |
| Listed as Toxic: YesNo                                 |
| Status of Environmental Compliance (Past and Present): |
|  |
|  |
|  |
| Known or Suspected Public Health Impacts:              |
| Reported to cause toxic hepatitis.                     |
|  |
| Known or Suspected Environmental Impacts:              |
|  |
|  |
|  |
| Level and Type of Worker Protection:                   |
| Lavel and Type of worker from the same                 |
|  |
| a di cara la Manga Information:                        |
| Source of Chemical Usage Information:                  |
|  |
| Source of Chemical Distribution Information/Accuracy:  |
|  |

LSTINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |   |
|--------------------------|---|
| 0ther                    |   |
| Offsite<br>Burial        |   |
| Onsite<br>Burial         |   |
| Onsite<br>Storage        | ·   |
| l.Iquid<br>Effluent      |   |
| Airborne<br>Release      | 00000   |
| Product                  | 5       6       5       6       5       6       5       6       6       6       6       6       6       6       6       6       7       6       7       6       7       7       6       7 |
| Quantity<br>Used<br>(Kg) | 91<br>91<br>182<br>273  |
| Year                     |   |
| Chemica 1                | Methylene<br>Dianiline (MDA)  |

| Chemical Name: Carbon Disulfide   |
|---|
| Uses:   |
| SolidLiquidXGas   |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):                              |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:   |
| Highly flammable, dangerous fire and explosion risk. Poisonous. Tolerance 20 ppm in |
|   |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection:  |
|   |
|   |
| Source of Chemical Usage Information:   |
| ·   |
|   |
| Source of Chemical Distribution Information/Accuracy:                               |
|   |

LSTINALLD CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                              | _ |
|--------------------------|------------------------------|---|
| 0ther                    |                              | _ |
| Offsite<br>Burial        | •                            | _ |
| Onsite<br>Burial         |                              | _ |
| Onsite<br>Storage        | •                            | _ |
| Liquid<br>Effluent       | •                            |   |
| Airborne<br>Release      | 1000%                        |   |
| Product                  |                              |   |
| Quantity<br>Used<br>(Kg) | 10.9<br>20.3<br>17.0<br>24.2 |   |
| Year                     | 1985<br>1984<br>1983<br>1982 |   |
| Chemical<br>Name         | Carbon Disulfide             |   |

| Chemica | .1 N | lame:  | Methylen    | e Chloride | <b>;</b>       |               |     |           |
|---------|------|--------|-------------|------------|----------------|---------------|-----|-----------|
| Uses:   |      |        |             |            |                |               |     |           |
| Solid_  |      | _ Liqu | id <u>X</u> | Gas        | <del></del>    | <del></del> · |     |           |
| Listed  | as   | Toxic: | Yes         | No         | <del>_</del> · |               |     |           |
| Status  | of   | Enviro | nmental     | Complia    | ance           | (Past         | and | Present): |
|         |      |        |             |            |                |               |     |           |

Known or Suspected Public Health Impacts: Chronic exposure causes loss of appetite, memory loss and disturbance of balance. It is an irritant to the respirator system and in high concentration causes pulmonary edema and pneumonitis, unconsciousned come and death. Acute oral exposure produces dizziness, drowsiness and unconsciousned It is an irritant to skin and eyes and in high concentration may cause chemical burns of the skin. The liquid has a defatting action on the skin. Acute skin absorption Known or Suspected Environmental Impacts: may cause the same effect as ingestion.

Level and Type of Worker Protection: Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respiratory protection.

Source of Chemical Usage Information:

LSTINALED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |   |
|--------------------------|---|
| Other                    |   |
| Offsite<br>Burial        |   |
| Onsite<br>Burial         |   |
| Onsite<br>Storage        |   |
| L. Iquid<br>Effluent     |   |
| Airborne<br>Release      |   |
| Product                  |   |
| Quantity<br>Used<br>(Kg) | 8,924<br>10,022<br>11,311<br>13,497<br>1,890<br>1,888 |
| Year                     | 1985<br>1984<br>1983<br>1980<br>1979                  |
| Chemical<br>Name         |   |

| Chemica | ll Name: | Acetonit | rile        |               |     |           |
|---------|----------|----------|-------------|---------------|-----|-----------|
| Uses:   |          |          |             |               |     |           |
| Solid_  | Lio      | quidX    | Gas         | <del></del> - |     |           |
| Listed  | as Toxio | : Yes    | No          |               |     |           |
| Status  | of Envis | onmenta  | l Compliand | e (Past       | and | Present): |

Known or Suspected Public Health Impacts:

Acute inhalation exposure can produce nose nad throat irritation, flushing, difficult breathing, chest pain, cyanosis, olfactory fatigue, cardiac and respiratory failure, coma and death. Oral exposure causes GI effects, difficult breathing, chest pain, irritation of mouth and throat, cyanosis and death. It can be absorbed through the skin and may cause similar effects. It is a skin and eye irritant.

Known or Suspected Environmental Impacts:

Biological effects have been reported in algae, protozoa, fathead minnows, bluegill and guppies.

Level and Type of Worker Protection:

Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber gloves, (3) safety glasses, (4) proper respirator protection.

Source of Chemical Usage Information:

# Estimated Distribution in %

| Unknown                  | 16<br>16<br>16<br>16   |
|--------------------------|--|
| 0ther                    |  |
| Offsite<br>Burial        |  |
| Onsite<br>Burial         |  |
| Onsite<br>Storage        |  |
| l.iquid<br>Effluent      |  |
| Airborne<br>Release      |  |
| Product                  |  |
| Quantity<br>Used<br>(Kg) | 19,757<br>24,329<br>21,063<br>14,369<br>10,776<br>15,512<br>15,675 |
| Year                     | LIGHMANICO   |
| Chemical<br>Name         | Acetonitrile   |

| Chemical Name: Niobium Pentachloride                                  |
|---|
| Uses:   |
| Solid_XLiquidGas  |
| Listed as Toxic: YesNo  |
| Status of Environmental Compliance (Past and Present):                |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts:                             |
| May evolve toxic fume of HCL. Possible kidney injury from small dose. |
| may evolve toxic rume of not. Tossible kraney induly from omer assets |
| Known or Suspected Environmental Impacts:                             |
| Maonin de daspososos  |
|   |
|   |
| Tours of Worker Protection:   |
| Level and Type of Worker Protection:                                  |
|   |
| * .   |
| Source of Chemical Usage Information:                                 |
|   |
| Source of Chemical Distribution Information/Accuracy:                 |

LSTINATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |  | ·       |   |
|--------------------------|--|---------|---|
| Other                    |  |         |   |
| Offsite<br>Burial        |  |         | · |
| Onsite<br>Burial         |  |         |   |
| Onsite<br>Storage        |  |         |   |
| Liquid<br>Effluent       |  |         |   |
| Airborne<br>Release      | *99<br>99<br>99<br>99<br>99                              |         |   |
| Product                  | 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4                  |         |   |
| Quantity<br>Used<br>(Kg) | 10,000<br>16,300<br>16,400<br>16,400<br>16,355<br>10,000 |         |   |
| Year                     | 1971<br>1970<br>1969<br>1968<br>1967<br>1966             |         |   |
| Chemica l<br>Name        | •  | *As HC1 |   |

| Chemical Name: Sodium Pentachlorophenate   |
|--|
| Uses:  |
| Solid Liquid Gas   |
| Listed as Toxic: Yes   |
| Status of Environmental Compliance (Past and Present):   |
|  |
| ··   |
|  |
| Known or Suspected Public Health Impacts:  |
| It is a skin, eye, and upper respiratory irritant.   |
|  |
| Known or Suspected Environmental Impacts:  |
| It is soluble in water and under laboratory conditions has shown acute toxicity to bluegill, channel catfish, rainbow trout and other aquatic organisms.   |
| ·  |
|  |
| Level and Type of Worker Protection:   |
| Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material includes the condition of the conditions of the condi |
| (1) adequate ventilation, (2) gloves, (3) safety glasses, (4) proper respiratory protection.   |
| Source of Chemical Usage Information:  |
|  |
|  |

# LSTINALLD CHEMICAL USAGE

Estimated Distribution in &

| Unknown                  |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             |             |
|--------------------------|--------|---------------------------------|--|--------|--------|--------|--------|--------|--------|---------------|--------|---|---|-------------|-------------|
| 0ther                    |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             |             |
| Offsite<br>Burial        |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             |             |
| Onsite<br>Burial         |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             |             |
| Onsite<br>Storage        |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             |             |
| t.iquid<br>Effluent      | 100%   | 100%                            | 100%                                     | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%          | 100%   | • |   |             |             |
| Airborne<br>Release      |        |                                 |  |        |        |        |        |        |        |               |        |   |   |             | <del></del> |
| Product                  | -      |                                 |  |        |        |        |        |        |        | <del></del> - |        |   |   |             |             |
| Quantity<br>Used<br>(Kg) | 28,000 | 28,000                          | 28,000                                   | 28,000 | 28,000 | 28,000 | 28,000 | 28,000 | 28,000 | 28,000        | 28,000 |   | - |             |             |
| Year                     |        | 1972                            | 1971                                     | 1970   | 1969   | 1968   | 1967   | 1966   | 1965   | 1964          | 1963   |   |   | <del></del> |             |
| Chemical<br>Name         |        | chlorophenate in<br>isopropanol | (Calgon Chem. Co.<br>or Nalco Chem. Co.) |        |        |        | •      |        |        |               |        |   |   |             |             |

| Chemical Name: Sodium Hydroxide   |
|---|
| Uses:   |
| Solid X Liquid Gas  |
| Listed as Toxic: Yes  |
| Status of Environmental Compliance (Past and Present):  |
|   |
|   |
|   |
| Known or Suspected Public Health Impacts: Chronic exposure to dusts may cause ulceration of nasal passages. Acute inhalation exposure causes upper respiratory tract irritation. Eye contact can result in damage to the cornea, conjunctiva and episclearal tissues. If skin is wet, edema and necrosis can occur. Ingestion can cause corrosion of tissue, vomiting, prostration, and collapse. |
| Known or Suspected Environmental Impacts:   |
|   |
|   |
|   |
| Level and Type of Worker Protection: Specific use, conditions, quantities and other factors determine the level and type of worker protection. Standard measures for protection against this material include: (1) adequate ventilation, (2) rubber clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.   |
| Source of Chemical Deage Information:   |

For the following chemicals, no descriptive data was located.

ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

|  | Unknown                  |  |
|--|--------------------------|--|
|  | Other                    |  |
|  | Offsite<br>Burial        |  |
|  | Onsite<br>Burial         | 2 2  |
|  | Onsite<br>Storage        | 000000   |
|  | Liquid<br>Effluent       |  |
| 1)                                     | Airborne<br>Release      |  |
|  | Product                  | 88888888<br>666666666  |
|  | Quantity<br>Used<br>(Kg) | 813<br>1,481<br>1,788<br>1,127<br>939<br>1,533<br>1,364<br>2,273 |
| •                                      | Year                     | 1985<br>1984<br>1982<br>1980<br>1970<br>1976<br>1976             |
| ************************************** | Chemical<br>Name         | Epoxy Resin (diglycidyl ether of bis phenol A)                   |

I STIMALLD CHEMICAL USAGE

Estimated Distribution in &

| <b>u</b> M               |  |  |      |  |             |
|--------------------------|--|--|------|--|-------------|
| Unknown                  |  |  |      |  |             |
| Other .                  |  |  |      |  |             |
| Offsite<br>Burial        | ,                                      |  |      |  |             |
| Onsite<br>Burial         | 86<br>86<br>86                         |  |      |  |             |
| Onsite<br>Storage        |  |  |      |  |             |
| Liquid<br>Effluent       | * ~ ~ ~                                |  |      |  |             |
| Airborne<br>Release      |  |  |      |  | <del></del> |
| Product                  |  | lution.                                    |      |  |             |
| Quantity<br>Used<br>(Kg) | 45,000<br>405,000<br>405,000<br>45,000 | nloride sc                                 | <br> |  | ·····       |
| Year                     |  | nium c                                     |      |  |             |
|                          |  | n ammo                                     |      |  |             |
| iom ica l                | Ammonium<br>Thiocyanate                | *Discharged in ammonium chloride solution. |      |  |             |

LSTIMALED CHEMICAL USAGE

|                             | Unknown                   |                                 |                                 |                                 |                       |           |      |   |      |         |   |
|-----------------------------|---------------------------|---------------------------------|---------------------------------|---------------------------------|-----------------------|-----------|------|---|------|---------|---|
| Ashpond &                   | Rogers<br>Quarry<br>Other | 50% to<br>pond only             | 20%                             | 80%                             | %06                   | %66       |      |   | <br> |         | _ |
| 4                           | Offsite<br>Burial         |                                 |                                 |                                 |                       |           |      |   |      |         |   |
| in 2                        | Onsite<br>Burial          |                                 |                                 |                                 |                       |           |      |   |      |         | _ |
| stribution                  | Onsite<br>Storage         |                                 |                                 |                                 |                       |           |      |   | <br> |         | _ |
| Estimated Distribution in 2 | t.iquid<br>Effluent       |                                 |                                 |                                 |                       |           | <br> |   |      |         |   |
| <u> </u>                    | Airborne<br>Release       | 20%                             | 20%                             | 20%                             | 10%                   | 1%        | -    |   |      | <u></u> | - |
|                             | Product                   |                                 |                                 |                                 |                       |           |      |   |      |         |   |
|                             | Quantity<br>Used<br>(Kg)  | 1955-19,200,000<br>1962 Kg/year | 1963-19,200,000<br>1967 Kg/year | 1962-19,200,000<br>1984 Kg/year | 19,200,000<br>Kg/year | 4,800,000 |      | • | <br> |         |   |
|                             | Year                      | 1955-<br>1962                   | 1963-<br>1967                   | 1962-<br>1984                   | 1985                  | 1986      |      |   |      |         |   |
|                             | Chemical                  | (Bottom and Fly)                |                                 |                                 |                       |           |      |   | •    |         | ٠ |

ESTIMATED CHEMICAL USAGE

Estimated Distribution in %

| Unknown                  |                      | , |       |      |   |
|--------------------------|----------------------|---|-------|------|---|
| 0ther                    |                      |   |       |      |   |
| Offsite<br>Burial        |                      |   |       |      |   |
| Onsite<br>Burial         | 10<br>10<br>20       |   |       |      |   |
| Onsite<br>Storage        |                      |   |       |      |   |
| Liquid<br>Effluent       |                      |   |       |      |   |
| Airborne<br>Release      |                      |   | <br>  |      | • |
| Product                  | 06<br>80             |   |       |      |   |
| Quantity<br>Used<br>(Kg) | 378<br>451<br>8629   |   | <br>- |      |   |
| Year                     | 1971<br>1970<br>1969 |   |       | <br> |   |
| Chemical<br>Name         | Pitch                |   |       |      |   |

LST INVILLE CHEMICAL, USAGE

Estimated Distribution in &

| Unknown                  |   |   |                                |                                |  |      |             |
|--------------------------|---|---|--------------------------------|--------------------------------|--|------|-------------|
| 0ther                    |   |   |                                |                                |  |      | <del></del> |
| Offsite<br>Burial        |   |   |                                |                                |  |      |             |
| Onsite<br>Nurial         | 70**<br>70                                |   |                                |                                |  |      |             |
| Onsite<br>Storage        |   |   | ate.                           |                                |  |      | <del></del> |
| l. Iquid<br>Effluent     | 30*                                       |   | sulfate and magnesium sulfate. |                                |  |      |             |
| Airborne<br>Release      |   |   | fate and ma                    | um sulfate.                    |  | <br> | <del></del> |
| Product                  |   |   |                                | fluoride and magnesium sulfate |  |      |             |
| Quantity<br>Used<br>(Kg) | 1,300,000                                 |   | acid sclution of calcium       |                                |  |      |             |
| Year                     |   |   | cid sc                         | nesium                         |  |      | <del></del> |
| Chemica]                 | Magnesium Fluoride/<br>Calcium Oxide Slag | • | *Discharged as weak            | **Calcium sulfate, magnesium   |  |      |             |

LSTINATED CHEMICAL USAGE

Estimated Distribution in &

| ite<br>al Other Unknown  |        | -      |        |        |        |        |        |        |        |        |        |              |   |      |
|--------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------------|---|------|
| Offsite<br>Burial        |        |        |        |        |        |        |        | -      |        |        |        |              |   |      |
| Onsite<br>Burial         |        |        |        |        |        |        |        |        |        |        |        |              |   | <br> |
| Onsite<br>Storage        |        |        |        |        |        |        |        |        |        |        |        |              |   | <br> |
| l.lquid<br>Effluent      | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   | 100%   |              |   | <br> |
| Airborne<br>Release      |        |        |        |        |        |        |        |        |        |        |        | ,            | - | <br> |
| Product                  |        |        |        |        |        |        |        |        |        |        |        |              |   |      |
| Quantity<br>Used<br>(Kg) | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | 19,000 | <del> </del> |   |      |
|                          |        |        |        | 0,     |        | 1968   | 1967   | 1966   | 1965   | 1964   | 1963   |              |   |      |
| Year                     | 1973   | 1972   | 1971   | 1970   | 1969   | 19(    | 19     | 119    | 119    | 139    | 119    |              |   | <br> |

### Y-12 HISTORICAL REVIEW URANIUM CONTAMINATED SCRAP HANDLING

Day 1 -- 1948
Scrap buried.

1948 to 1955

Scrap was stored in White Wing holding yard. The material was sold to Witherspoon.

1955 to present

Scrap taken to west end of Y-12 and stored.

The material was stored at Y-12 from 1955 until Oak Ridge Processing Company, Inc. started smelter operations. The smelter processed copper starting February, 1957 and started steel smelter in August 1957. The Company smelted contaminated scrap accumulated from 1955 until the smelter closed down in May 1960.

Witherspoon began purchase of the contaminated scrap in September 1963 for remelt at Knoxville Iron Smelter Plant. This contract was terminated December 1964, after a total of 2,856 tons was purchased.

Scrap was then accumulated in the yards and sold as follows:

| Date            | Tonnage | Purchaser   |
|-----------------|---------|-------------|
| March, 1968     | 440     | D. D. Smith |
| September, 1968 | 550     | Witherspoon |
| April, 1970     | 782     | Witherspoon |
| June, 1972      | 751     | D. D. Smith |

### SCRAP SALES

| 9/63-2/64        | 2005 G.T. U Contaminated Scrap                    | David Witherspoon                  |
|------------------|---|------------------------------------|
| 2/64-12/64 (719) | 851 G.T. U Contaminated Scrap                     | David Witherspoon                  |
|                  | 8000# Beryllium Scrap                             | Unknown                            |
| 7/15/64 (743)    | 23,000# Nickel Scrap                              | Lee Metals Corp.                   |
| 10/74 (754)      | 23,000# NICKEL Borah                              | Twin City Iron and                 |
|                  | 32,780# Lead Covered Cable                        | Metal Co.                          |
| 2/23/65 (773)    | 12,500# Lead                                      | Karl Nussbaum and Son, Inc.        |
|                  | 7,500# Nickel                                     | Lee Metals Corp.                   |
| 3/9/65 (777)     | Nickel Scrap                                      | Allied Metal Co.                   |
| 3/9/05 (111)     | Monel Scrap                                       | Allied Metal Co.                   |
| 1. (00/65 /F94)  | 6,017# Beryllium Chips                            | Unknown                            |
| 4/28/65 (781)    | 44 COOA Ethelana Dibromida                        | Unknown                            |
| 5/10/65 (782)    | 11,700# Ethylene Dibromide                        | Twin City Iron and                 |
| 8/19/65 (800)    | Lead  | Metal Company, Inc.                |
|                  |   |                                    |
|                  | Lead Covered Cable                                | Brodey & Brodey, Inc.              |
| 10/8/65 (817)    | 19,489# Beryllium Chips and                       | Unknown                            |
|                  | Powder  |                                    |
| 12/6/65 (822)    | 2.100# Lead Covered Cable                         | David Witherspoon                  |
|                  | 58,000# Nickel Plate, Hg. Contam.                 | Samuel G. Keywell,                 |
|                  | , ,   | Company, Inc.                      |
|                  | 223,700# Mixed Metal, Hg. Contam.                 | -                                  |
|                  | 613,160# Scrap Metal, Hg. Contam.                 |                                    |
|                  | 59,500# Nickel, Hg. Contam.                       | Samuel G. Keywell,                 |
|                  | 99,900F HTCZCZ, HB. 00H1                          | Company, Inc.                      |
|                  | 153,700# Structural Steel,                        | David Witherspoon                  |
|                  | = <del>•</del>                                    | Baraa na saas op saas              |
|                  | Hg. Contam.                                       | Knoxville Junk Co.                 |
|                  | 32,000# Pipe, Valves, etc.,                       | MOXVIIIE OMIA 00.                  |
|                  | Hg. Contam.                                       | Toward 11 o Tumbe Co               |
| 1/14/66 (826)    | 2,108,000# Mixed Metal,                           | Knoxville Junk Co.                 |
|                  | Hg. Contam.                                       | - 1 1 G. T                         |
|                  | 96,600# Nickel Scrap, Hg. Contam.                 | Frankel Co., Inc.                  |
| 2/9/66 (828)     | 11,700# Ethylene Dibromide                        | Unknown                            |
| 4/26/66 (840)    | 18,032# Beryllium Chips & Powder                  | Unknown                            |
| 5/13/66 (841)    | 5,100# Lead Encased in S.S.                       | David Witherspoon                  |
|                  | 87,000# Al. Scrap, Hg. Contam.                    | M. Kimerling and Sons, Inc.        |
|                  | 92,500# Stainless Steel Scrap,                    | Samuel G. Keywell                  |
|                  | Hg. Conta.  | Company, Inc.                      |
|                  | 21,500# Bare Brass Scrap,                         | M. Kimerling and                   |
|                  | *   | Sons, Inc.                         |
|                  | Hg. Contam.<br>104,000# Nickel Scrap, Hg. Contam. |                                    |
|                  |   | Stainless Processing               |
|                  | 101,000# Nickel Plates,                           |                                    |
|                  | Hg. Contam.                                       | Company<br>Karl Nussbaum and       |
|                  | 280,800# Scrap Pipe, Hg. Contam.                  |                                    |
|                  |   | Sons, Inc.                         |
|                  | 25,000# Scrap Instruments,                        | Twin City Iron and                 |
|                  | Hg. Contam.                                       | Metal Co., Inc.                    |
|                  | 19,500# Lead Scrap                                | Twin City Iron and Metal Co., Inc. |
|                  | 178,603# Mixed Metal Scrap,                       | Pits Scrap Iron and                |
|                  | Hg. Contam.                                       | Metal Company                      |
| 7/6/66 (849)     | 209,000# Scrap Nickel Plate,                      | Lee Metals Corp.                   |
| 1,0,00 (073)     | Hg. Contam.                                       | -                                  |
|                  | 2,000,000# Mixed Metals,                          | Hunsaker Iron and                  |
|                  | Hg. Contam.                                       | Metal Company                      |
|                  | *0. ***********************************           | •                                  |

Lee Metals Corp. 67,500# Copper Scrap, Hg. Contam. Richelson Iron and 43,500# Copper Scrap, Hg. Contam. Metal Company David Witherspoon 12,500# Lead Covered Cable 2/7/67 (874) Berman Brothers Iron 74.000# Al Bus Bar, Hg. Contam. & Metal Co., Inc. Metropolitan Metals, 99,000# Al Scrap, Hg. Contam. Commercial-London, 97,000# Bare Copper Scrap, Inc. Hg. Contam. David Witherspoon 37,000# Insulated Copper Scrap, Hg. Contam. 21,000# Mixed Brass & Bronze Scrap, Chicago Metals Corp. Hg. Contam. U.S. Reduction Co. 4,300# Nickel Scrap, Hg. Contam. Wolverine Metal Co. 133,000# Stainless Steel Scrap, Hg. Contam. McGean Chemical Co. 1 Lot Depleted Uranium Oxide 4/3/67 (882) 40,320# Depleted Uranium Oxide Unknown 4/27/67 (883) Rockwood Iron and 2,700,000# Mixed Scrap Metal, 10/24/67 (910) Metal Company Hg. Contam. Richelson Iron and 137,000# Scrap Pipe, Hg. Contam. Metal Company Knoxville Junk Co. 67.000# Bare Copper Scrap, Hg. Contam. Lee Metals Corp. 120,000# Aluminum Scrap, Hg. Contam. Lee Metals Corp. 61.000# Copper Scrap, Hg. Contam. Lee Metals Corp. 28.000# Bare Brass Scrap, Hg. Contam. David Witherspoon 11.000# Mixed Copper & Ferrous Scrap. Hg. Contam. 36,000# Aluminum Bus., Hg. Contam. Berman Brothers Iron and Metal Wolverine Metal Co. 2,070# Nickel Scrap Wolverine Metal Co. 3,540# Monel Scrap David Witherspoon 21.000# Lead Scrap 40,200# Depleted Uranium Oxide Unknown 11/21/67 (921) Brush Beryllium Co. 5,896# Beryllium Chips 2/27/68 (932) The Beryllium Corp. 3,007# Beryllium Chips David Witherspoon 600 G.T. U Contaminated Scrap 3/5/68 (933) Unknown 402 G.T. U Contaminated Mixed Metal Scrap 40,200# Depleted Uranium Oxide McGean Chemical Co. 4/2/68 (942) David Witherspoon 1 Lot Lead Covered Cable 4/17/68 (946) Richelson Iron and 73.000# Insulated Copper Scrap, Metal Company Hg. Contam. M. Kimerling and 40,000# Bare Copper Scrap, Sons, Inc. Hg. Contam. David Witherspoon 105,000# Aluminum Scrap, Hg. Contam. Mercer Alloys Corp. 175,000# Stainless Steel Scrap, Hg. Contam.

| 6/12/68 (956)<br>8/1/68 (964)<br>8/13/68 (967)<br>9/17/68 (984) | 16,038g. Rare Earths 39,600# Depleted Uranium Oxide 3,239# Beryllium Oxide 851,000# U Contaminated Carbides 134,000# U Contaminated Diffusion Pumps | Research Chemical McGean Chemical Co. Brush Beryllium Co. David Witherspoon David Witherspoon |
|---|---|---|
|   | 77,400# U Contaminated Trans-<br>formers and Rectifiers<br>80,000# U Contaminated Dollies   | David Witherspoon  David Witherspoon  |
|   | and Jacks   | •   |
|   | 90,000# U Contaminated Scrap Metal  | David Witherspoon   |
| 10/3/68 (989)   | 289# Niobium Pentachloride  | Unknown   |
| 11/7/68 (991)   | 1,360,000# Mixed Scrap Metal,   | Knoxville Junk Co.  |
|   | Hg. Contam.<br>3,190# Nickel Scrap  | Wolverine Metal   |
|   |   | Company, Inc.   |
|   | 11,800# Lead Covered Cable Scrap  | David Witherspoon   |
| 1/21/69 (999)   | 40,200# Depleted Uranium Oxide  | McGean Chemical Co.   |
| 3/19/69 (1006)  | 39,600# Depleted Uranium Oxide  | McGean Chemical Co.<br>McGean Chemical Co.  |
| 5/8/69 (1012)   | 80,400# Depleted Uranium Oxide  | Lee Metal Corp.   |
| 5/13/69 (1013)  | 12,900# Lead Covered Cable  | East Tennessee Iron   |
|   | 644.4 G.T. Mixed Scrap Metal,<br>Hg. Contam.  | and Metal Co.   |
| 9/23/69 (1030)  | 951# Niobium Pentachloride  | Alfa Inorganics, Inc.   |
| 10/1/69 (1033)  | 2,930,000# Mixed Scrap Metal,   | Rockwood Iron and   |
| 107 17 0 7 (1033)   | Hg. Contam.   | Metal Company   |
| 10/15/69 (1035)   | 380# Beryllium Compounds  | Unknown   |
| 10/22/69 (1040)   | 40,200# Depleted Uranium Oxide  | Chemetron Corporation   |
| 12/10/69 (1042)   | 1,760,910# Mixed Scrap Metal,   | Rockwood Iron and   |
|   | Hg. Contam.   | lverine-Sitkin  |
|   | 2,800# Nickel Scrap   | Alloys, Inc.  |
|   | 4,300# Lead Covered Cable Scrap   | Knox Metals Corp.   |
|   | 14,500# Lead Scrap  | Lee Metals Corp.  |
|   | 1,700# Monel Scrap  | Wolverine-Sitkin  |
|   |   | Alloys, Inc.  |
| 2/18/70 (1048)  | 1 Lot Nitric Acid   | David Witherspoon   |
| 2/26/70 (1056)  | 9,675# Nickel Ingots  | H. Klaff and Company  |
| 3/24/70 (1062)  | 2 Lots Nickel Ingots  | Mercer Alloys Corp.   |
|   | 39,269# Nickel Ingots   | Lee Metals Corp. Chemetron Corp.  |
| 4/16/70 (1069)  | 40,200# Depleted Uranium Oxide  | David Witherspoon   |
| 5/12/70 (1072)  | 600 G.T. U Contaminated Scrap Metal   | PSAIR WIREL Shoot   |
| 9/15/70 (1088)  | 500# U Contaminated Scrap Metal   | David Witherspoon   |
| 1/27/71 (1095)  | 1 Lot Nickel Sulfamate  | DuPont Smith  |
| 11/16/71 (2047)   | 200 G.T. U Contaminated Scrap Metal   | I. Deutch & Sons  |
|   | 3,000# Monel Scrap  | Commercial Metals Company   |
| 2/23/72 (2064)  | 6,000# Nickel Scrap   | S. G. Keywell,  |
| •   | •   | Company, Inc.   |
|   | 1,675# Monel Scrap  | Tecmet, Inc.  |

|                 | 3,000# Lead Covered Copper Scrap<br>46,500# Lead Scrap<br>1,520# Zirconium                            | Knox Metals, Corp.<br>Lee Metals, Corp.<br>Wolverine-Sitkin<br>Alloys, Inc. |
|-----------------|---|---|
| 5/17/72 (2085)  | 6 Lots Mercury Contaminated Material  | Mallory Battery Co.   |
| 7/18/72 (2100)  | 700 G.T. U Contaminated Mixed Metal   | DuPont Smith  |
| 8/21/73 (2134)  | 21,080# Lead Scrap<br>19,700# Nickel Scrap<br>2,620# Lead Covered Cable Scrap<br>520# Zirconium Scrap | Unknown<br>Unknown<br>Unknown<br>Unknown                                    |
| 7/17/74 (2181)  | 890# Nickel Scrap<br>29,280# Lead Covered Cable Scrap   | Commercial Metals Co.<br>Unknown  |
| 8/1/74 (2177)   | 23 Lots Scrap Beryllium   | Multiple Buyers   |
| 8/26/75 (2232)  | 4,600# Lead Scrap   | Unknown   |
| 7/13/77 (2302)  | 3,070# Monel Scrap  | Unknown   |
| 3/22/79 (2405)  | 440 Gallons Benzene   | Unknown   |
| 1/29/80 (2451)  | 10 Lots Beryllium   | Unknown   |
| 12/10/81 (2575) | 1 Lo Lead Scrap   | David Witherspoon   |


.

•



Mailed\_

| (This section to be completed by ChemRisk)   |
|--|
|  |
| Sandberg / CEP  Name Division is requested to provide the following document                       |
| Name Division is requested to provide the following document                                       |
| Address  |
| Date of Request 1/25/93 Expected receipt of document   |
| Title of requested document Historic Chemical Release Report for                                   |
| Dak Rine National Laboratory (ORNL)  |
|  |
| Document Number Unnumbered 16-03752  |
| Access Number of Document Date of Document 4-25-86   |
|  |
| (This section to be completed by Derivative Classifier)  Derivative Classifier August Phone 7-7864 |
| Date document transmitted to Dr. Quist   |
| Date release received from Dr. Quist   |
| PUBLIC RELEASE STAMP attached to each copy of document (YES NO)                                    |
| Date document sent to reproduction Expected Return   |
| Delivered to DRC byDate  |
|  |
| (This section to be completed by DRC)  |
| Received by DRCDate  |
| Processed  |



# OAK RIDGE K-25 SITE DOCUMENT RELEASE FORM

| Person requesting release MAMAS E Widner Chem 7 . Telephone No. 4-6745   |
|--|
| Person requesting release 7/1/1/100 WIGHT WIGHT WILL 129193 — Telephone No. 9-0793   |
| Mailing Address K-1/00 W/3 +262 11 251 ision or Organization   |
| Person requesting release MMM Widner (William) Telephone No. 9-079  Mailing Address K-1200 M5 7262 1 29 93 — Telephone No. 9-079  Date by which release is required Some documen Recid K-2500-RC (4710) ison or Organization  Some documen Recid K-2500-RC (4710) g time will be longer).  Note: Two copies of the document must generally . AS Quist, and Information Control Office with this request. Only  |
| Note: Two copies of the document must generally . AS Quitarian and Information Control Office with this request. Only one copy of photos and videotapes is required an include photos must be accompanied by "originals" of the photos.  |
| Approval of request for Classification and Information Control Office to release document (department head or higher):  Signature: Date  |
| DOCUMENT DESCRIPTION (to be completed by requester)  |
| Document number LANNumbered 603752  Document title Historic (Nemical Release Report for Oak Ridge National Laboratory (ORNL)  Author(s) (Indicate other divisions or organizations, if applicable)  Pages  Pa |
| Document type (See Doc. Prep. Guide, Chs. 1 and 2, for definitions of document types):   |
| ☐ Formal Report ☐ Progress Report ☐ Informal R&D Report ☐ Abstract ☐ Drawing   |
| ☐ Administrative ☐ Correspondence ☐ Internal Technical Data ☐ Photo ☐ Other Visuals  |
| Journal Article (identify journal):  |
| Oral Presentation (identify meeting, sponsor, location, date):   |
| Will oral presentation be published in program, booklet, brochure, etc.?   |
| Is copyrighted material contained in this document? (If present, attach release.)  |
| Remarks  |
| CLASSIFICATION INFORMATION (to be obtained by requester)   |
| Was the work reported in this document funded, in whole or in part, by a classified program at Martin Marietta Energy Systems, Inc.?    No   |
| Is the subject area of this document closely related to a prior or current classified program at Martin Marietta Energy Systems, Inc.?   |
| ☐ No ☐ Yes Within the Department of Energy? ☐ No ☐ Yes   |
| Name or Description of applicable program(s)   |
|  |
| Additional remarks   |
| This document contains no classified information./ / / 1   |
| Derivative Classifier signature and wilch Date 1/27/93   |
| UCN-87 (1 7-91)  |

| DISTRIBUTION LIMITATIONS (If any) (completed by requester) |                    |                              |                     |                                       |
|--|--------------------|------------------------------|---------------------|---------------------------------------|
| ☐ Unrestricted, unlimited                                  |                    |                              |                     |                                       |
| Distribution may be limited because this document con      | ntains information | n that is:                   |                     |                                       |
| ☐ Unclassified Controlled Nuclear Information *            | ☐ Applied 7        | echnology *                  |                     | Export Controlled *                   |
| ☐ Naval Nuclear Propulsion Information *                   | ☐ Gov't Co         | fidential Commercial Inform  | mation *            | ] Proprietary                         |
| ☐ Sensitive Nuclear Technology *                           | ☐ Small Bu         | siness Innovation Research   | .* [                | Official Use Only                     |
| ☐ Safeguards Information *                                 | ☐ Cooperat         | ve R&D Agreement *           |                     | ] Other                               |
| -  | * Ge               | nerally identified by sponso | r                   |                                       |
|  |                    |                              |                     |                                       |
| Remarks:   |                    |                              |                     |                                       |
| Tionano.   |                    | <del></del>                  |                     |                                       |
|  |                    |                              |                     |                                       |
| PATENT IN  | FORMATION (        | completed by requester)      |                     |                                       |
| •  | •                  | <u></u>                      |                     |                                       |
| Does this document disclose any new equipment, pro         | cess, or materia   | 1?                           | ☐ No                |                                       |
| If yes, list the patent significance and identify page nur | mber(s) and line   | number(s) in the space in    | nmediately followin | g                                     |
| (or attach separate pages).                                |                    |                              |                     |                                       |
|  |                    | <del></del>                  | ····                | · · · · · · · · · · · · · · · · · · · |
| PATENT SECTION ACTION (completed by Pater                  | nt Section upo     | request by the Classific     | ation and Informa   | tion Control Office)                  |
| ./   |                    |                              |                     | · · · · · · · · · · · · · · · · · · · |
| Document may be released for publication                   |                    | Oocument must be reviewe     | -                   | aroup before release                  |
| Document contains pate                                     | entable informa    | ion and may not be release   | ed at this time     |                                       |
| Remarks  |                    |                              |                     |                                       |
| Patent Section Representative                              |                    | Dat                          | e $2/2/$            | 193                                   |
| ratein Section Representative                              |                    | - Jan                        |                     |                                       |
| CLASSIFICATION AND INFORMATION CONTROL                     | OFFICE ACTI        | N (completed by Classifi     | cation and inform   | ration Control Office)                |
| CEASSIFICATION AND INFORMATION CONTROL                     | OFFICE ACTI        | on (completed by Classin     | cation and inform   | ladon Control Office)                 |
| Classification Office                                      | •                  | ☐ Approved for a             | elease with chang   | es (see below)                        |
| Action Taken:  Approved for release with                   | hout change        |                              |                     |                                       |
|  |                    |                              |                     |                                       |
|  |                    |                              |                     |                                       |
| Classification Officer signature                           | 82                 | Da                           | te 2/2/             | 9)                                    |
| <u> </u>   |                    |                              |                     |                                       |
|  |                    |                              | <del></del>         |                                       |
| Technical Information  Not approved for releas             | e (see below)      | ☐ Approved for               | release with change | ges (see below)                       |
| Office Action Taken: Approved for release wi               |                    |                              |                     |                                       |
| /  | •                  | •                            |                     |                                       |
|  |                    |                              |                     |                                       |
|  | <del></del>        | <del></del>                  |                     | ·                                     |
|  | 79 (               |                              |                     |                                       |
| Technical Information Officer Signature                    | X de               | est Dai                      | e 2/2/              | 93                                    |
|  |                    | , Dai                        |                     |                                       |
| Send to OSTI? Yes No                                       |                    | Category Distribution:       |                     | <del> </del>                          |

**DOCUMENT NUMBER:** UNNUMBERED/G03752

DOCUMENT TITLE: HISTORICAL CHEMICAL LISTING - ORNL

**AUTHORS:** 

**DOCUMENT TYPE:** ATTACHMENT 1

**DOCUMENT DATE:** 

PURPOSE OF RELEASE: HEALTH STUDY FEASIBILITY REPORT

COPY RIGHTED MATERIAL: NO

**DOCUMENT NUMBER:** UNNUMBERED/G03752

**DOCUMENT TITLE: ORNL SURPLUS CHEMICAL SALES AND TRANSFERS** 

**AUTHORS:** 

**DOCUMENT TYPE: ATTACHMENT 2** 

**DOCUMENT DATE:** 

PURPOSE OF RELEASE: HEALTH STUDY FEASIBILITY REPORT

COPY RIGHTED MATERIAL: NO



KR

MARTIN MARIETTA ENERGY SYSTEMS, INC.

April 25, 1986

C. G. Jones, 1000, 211A, MS 214A (4-9969)

Historic Chemical Release Report for Oak Ridge National Laboratory (ORNL)

Reference: Correspondence from R. L. Egli to W. F. Furth, "Historic Chemical Release Report," dated January 6, 1986

As I indicated to you earlier, we have not been able to complete the subject report for ORNL by the requested deadline of April 25. Historically, the research activities of the Laboratory have utilized numerous chemicals. Because of this wide array of chemicals, it has proven to be a time consuming task to sort through available records in order to identify the historic use and disposition of toxic chemicals with the greatest potential for environmental or public health impacts, as well as the large quantity materials.

At this time we are providing a portion of the requested information. Enclosed is a listing of chemicals and their quantities which have been identified to be included in the ORNL report (Attachment 1). A Chemical Release Data Form and the Estimated Chemical Usage Form will be completed for each of the chemicals on this list. Also enclosed is a summation of ORNL surplus chemical sales and other related activities (Attachment 2).

Our current plans are to provide the completed forms referenced above, and other pertinent information on or before May 9, 1986. I hope this delay does not cause any inconvenience to you. If there are any questions, please call.

P. S. Rohwer, 4500S, G-260, ORNL (4-6670)

Rohwer/an

PSR:BME:871N:aw

cc/enc: B. M. Eisenhower

D. C. Parzyck

APPROVAL FOR RELEASE

Unnumbered 1-page 1tr, PS Rohwer to CG
Document: # Jones "Historic, Chemical Release Report for ORNL; dtd 4/25/86; 16
pages attachment.

Approval for unrestricted release of this document is authorized by the Oak Ridge K-25 Site Classification and Information Control Office, Martin Marietta Energy Systems, Inc., PO Box 2003, Oak Ridge, TN 37831-7307.

K-25 Classification & Information Control Officer

Date

#### ATTACHMENT 1

Historical Chemical Listing - ORNL

# DRAET

| Chemical                       | Quantit | <u>y*</u> |
|--------------------------------|---------|-----------|
| Acetone                        | 83,789  | L         |
| Acetonitrile                   | 135,594 |           |
| Acrylamide                     | 1,389   |           |
| Aluminum nitrate               | 7,455   | -         |
| Ammonia, anhydrous             | 20,580  | _         |
| Ammonium bifluoride            | 657     | -         |
| Ammonium hydroxide             | 8,800   | _         |
| Ammonium nitrate               | 4,445   | kg        |
| Asbestos-Containing Materials  |         |           |
| Asbestos Felt Roofing          | 8,627   |           |
| Asbestos Sheet Roofing         | 6,399   |           |
| Asbestos Tape                  | 246     | rolls     |
| Asbestos Insulated Wire        | 1,649   | m         |
| Asbestos Gaskets               | 17,453  |           |
| Asbestos Packing               | 1,838   |           |
|                                | 971     |           |
| Barium octahydrate             | 1,918   |           |
| Baygon (N-Methyl carbamate)    | 15      | L         |
| Benzene                        | 2,589   |           |
| Cadmium nitrate                | 183     |           |
| Calcium hypochlorite           | 7,297   |           |
| Carbon monoxide                | 267     |           |
| Carbon tetrachloride           | 1,499   |           |
| Chlordane                      | 189     |           |
| Chlorine, gas                  | 9,692   | _         |
| Chloroform                     | 1,067   |           |
| Chlorpyrifos (Bolt)            | 136     |           |
| Chromic acid                   | 8,482   |           |
| Coal-oil mixture               | 18,925  |           |
| Coal Tar Pitch Cyclohexane     | 22,014  |           |
| Diazinon                       | 386     |           |
| 2,4—Dichloropenoxy acetic acid | 19      |           |
| Diethyl benzene                | 189     |           |
| Diethylenetriamine pentacetic  | 2,185   | кд        |
| sodium salt                    | 545     | ka        |
| 1,4-Dioxane                    | 155     | _         |
| Disulfoton (Disyston G)        |         | kg        |
| Dodecane                       | 3,142   | _         |
| Endcor (Mixture of sodium salt | 0,112   | _         |
| organo-phosphoric acid and     |         |           |
| Merceptobenzotriazolo)         | 10,825  | L         |
| Epoxy Resins                   | 447     |           |
| Ethyl acetate                  | 1,787   |           |
| Ethyl alcohol                  | 46,749  |           |
| Ethyl alcohol, denatured       | 15,083  | L         |
| Ethyl ether                    | 3,064   | kg        |
| Ethylene glycol                | 114,421 | L         |
| Ethylene oxide                 | 100     |           |
| Ferric chloride                | 4,924   |           |
| Ferric sulfate                 | 1,000   | kg        |
| Formaldehyde                   | 1,177   |           |
| Formamide                      | 469     | L         |

# DRAET

| Chemical                                   | <u>Quantity*</u>    |
|--|---------------------|
| Hexane                                     | 6,491 L             |
| Hydrochloric acid                          | 2,633,145 L         |
| Hydrofluoric acid                          | 6,884 kg            |
| Hydrogen peroxide                          | 12,463 kg           |
| Isopentane                                 | 409 L               |
| Isopropyl alcohol                          | 79,228 L            |
| Kelthane (4,4'-Dichloro-cx-trichloromethyl | •                   |
| benzhydrol)                                | 19 L                |
| Lacquer thinner                            | 48,770 L            |
| Lasso (2—Chloro-2'6'—diethyl—N—            |                     |
| Methoxymethyl acetanilide                  | 9 L                 |
| Lead, sheet                                | 30,551 kg           |
| Lindane                                    | < 1 kg              |
| Magnesium nitrate                          | 830 kg              |
| Malathion                                  | 242 L               |
| Mercury                                    | 1,687 kg            |
| Methyl alcohol                             | 843,199 L           |
| Methyl ethyl kelone                        | 10,008 L            |
| p,p' Diaminodiphenylmethane                | 91 kg               |
| Methylene chloride                         | 67,403 L            |
| Metex Stripper                             | 3,345 kg            |
| Micro-Bio-Treat (Mixture of n-alkyl        |                     |
| dimethyl benzyl ammonium                   |                     |
| chloride and Bis [Tri-n-                   |                     |
| butyltin] oxide)                           | 5,072 L             |
| Naphtha                                    | 24,565 L            |
| Nickel chloride                            | 4,805 kg            |
| Nickel sulfate                             | 2,590 kg            |
| Nitric acid                                | 112,191 L           |
| Nitric oxide                               | 20 m <sup>3</sup>   |
| Orthene (Acetylphosphoramidothioic acid -  |                     |
| 0,5—dimethyl ester)<br>Pararosaniline      | 4 kg                |
| Pentane                                    | 83 L                |
| Perchloric acid                            | 382 L               |
| Petroleum ether                            | 2,017 kg<br>1,506 L |
| Phenol                                     | 242 kg              |
| Phenolic resin                             | 242 kg<br>91 kg     |
| Phosphoric acid                            | 9,190 kg            |
| Polychlorinated biphenyls                  | 9,190 kg            |
| Potassium cyanide                          | 200 kg              |
| Potassium hydroxide                        | 557,677 kg          |
| Pramitol                                   | 277 kg              |
| Pyridine                                   | 159 L               |
| Ronstar                                    | 455 kg              |
| Round-up (N-Phosphonomethyl glycine        |                     |
| isopropylamine salt)                       | 231 L               |
| Silicon tetrafluoride                      | 3,125 kg            |
| Silvex                                     | 284 L               |
| Sodium bifluoride                          | 6,455 kg            |
| Sodium cyanide                             | 2,370 kg            |
| Sodium dithionite                          | 807 kg              |
| Sodium hexametaphosphate                   | 455 kg              |
|  |                     |

| Chemical                             | Quantity* |    |
|--------------------------------------|-----------|----|
| Sodium hydroxide                     | 394,646   | kq |
| Sodium nitrate                       | 70,256    | _  |
| Sodium sulfide                       | 500       | _  |
| Subdue E2                            | 4         | L  |
| Sulfur hexafluoride                  | 7,214     | kq |
| Sulfuric acid                        | 45,965    | L. |
| Talon (4'-Bromo (1,1'-biphenyl-4-yl) | 68        | ka |
| Tetrachloroethylene                  | 2,147,321 | _  |
| Tetrahydrofuran                      | 2,010     |    |
| Toluene                              | 9,281     |    |
| Tributyl phosphate                   | 5,934     | kq |
| Trichloroethylene                    | 49,148    | L  |
| Tichloromethylsilane                 | 57        |    |
| Triethanolamine                      | 92        |    |
| Urethane                             | 984       | _  |
| Varsol                               | 181,218   | L  |
| Vydate                               | 23        | L  |
| Xylene                               | 1,692     | L  |
|                                      | 54        | kg |
|                                      |           |    |

\*Quantities were determined from the survey of purchase requisitions for chemical purchases and ORNL Chemical Stores issues using ORNL account numbers.

#### ATTACHMENT 2

ORNL Surplus Chemical Sales and Transfers

## ORNL Waste Oil Sold for Recycle

| <u>Date</u>        | Quantity (liters) | Off-site Buyer  |
|--------------------|-------------------|---|
| FY 1980            | 37,850            | Petroleum Recycling Corp.                                 |
| FY 1981<br>FY 1982 | 39,364<br>25,/38  | Alpha Recovery Systems, Inc.<br>Petroleum Recycling Corp. |
| FY 1983            | 8,176             | Universal Oil & Supply Co.                                |
| FY 1984<br>FY 1985 | 71,468<br>7.949   | Alpha Recovery Systems, Inc. Universal Oil & Supply Inc.  |
|                    | 18,168            | Alpha Recovery Systems, Inc.                              |
| Total              | 208,713           |   |

All wastes oils are analyzed for PCBs, organics, and radioactivity prior to being placed in the Waste Oil Storage Tank.  $\cdot$ 

#### Waste Acceptance Criteria:

| PCB              | <2 ppm                     |
|------------------|----------------------------|
| Organics         | <50 ppm                    |
| Gross beta       | $1.1 \times 10^{-2}$ Bq/mL |
| Gross gamma      | $1.1 \times 10^{-2}$ Bq/mL |
| 3 <sub>H</sub> - | 1.1x10 <sup>3</sup> Bq/mL  |
| 14 <sub>C</sub>  | $3 \times 10^2$ Bg/mL      |



## Surplus Chemicals Transferred to Other Government Agencies

| <u>Date</u> | Chemical  | <u>Quantity</u>  | Receiving Agency                          |
|-------------|---|--|---|
| 6/79        | Iodic acid Nickelous sulfate Potassium chromate Potassium permanganate Hardening resin Argon  | 1 kg<br>4 kg<br>9 kg<br>23 kg<br>57 g<br>6 L                           | Department of General<br>Services         |
| 6/80        | 1-Amino-8-naphthol-4- sulfonic acid Benzoic anhydride acid 1-Naphthylamine-4- sulfonic acid 1-Naphthylamine-8- sulfonic acid Benzyl acetate 2,4-Dichloro-1-Naphthol 2,3-Dichloronaphthoquinone 2,7-Dihydroxynaphthalene A-Fluoronaphthalene A-Naphthonitrile B-Naphthylamine hydrochloride A-Naphthylhydrazine hydrochloride 6-Nitroquinoline   | 1000 g 100 g 25 g 1 kg 2 kg 100 g 25 g 50 g 10 g 50 g 10 g 50 g 40 g   | Comparative Animal<br>Research Laboratory |
| 7/80        | o-Cresotinic acid 2,4-Dinitro-1-naphthol-7- sulfonic acid P-Fluorobenzoic acid 8-Hydroxy-5-quinoline sulfonic acid 7-Iodo-8-hydroxyquinoline- 5-sulfonic acid A-Naphthaleneacetic acid B-Naphthoxyacetic acid S-Nitrobarbituric acid S-Oxynaphtholic acid Quinaldinic acid Violuric acid 1-Amino-2-naphthol hydrochloride 2-Aminonaphthoquinone 2-Chloroquinoline 6-Chloroquinoline 5,7-Dibromo-8- hydroxyquinoline | 100 g 25 g 5 g 100 g 100 g 100 g 25 | Comparative Animal<br>Research Laboratory |

# ORAFT

## Surplus Chemicals Transferred to Other Government Agencies

| Date          | Chemical   | Quant               | <u>tity</u>                           | Receiving Agency  |
|---------------|--|---------------------|---------------------------------------|---|
| 7/80<br>cont. | 2,4-Dibromo-1-naphthol 9,10-Dichloroanthracene 3,5-Dimethylpyrozole 2,7-Dinitrophenanthra- quinone 2,4-Dioxothiazolidine N-Methyl-2-quinoline 5-Nitro-1-naphthylamine 2-Nitrophenanthraquinone | 25<br>25<br>5       | g<br>g<br>g                           |   |
|               | 8-Nitroquinoline   | 10                  | -                                     |   |
| 8/80          | 1 lot (507 items) of miscellaneous chemicals classified as DOT Flammab liquid, n.o.s. and poisonous solid, n.o.s.  | le                  |                                       | University of Tennessee,<br>Knoxville via the<br>Department of General<br>Services  |
| 7/83          | Potassium nitrite  | 16                  | kg                                    | Georgia Coastal Plain<br>Experimental Station,<br>Tifton, GA                        |
| 12/85         | Ethyl Alcohol  | 11                  | drums                                 | University of Tennessee,<br>Knoxville via the<br>Departament of General<br>Services |
| 3/86          | Hydraulic fluid,<br>non-petroleum base<br>Hydraulic fluid<br>Gulf, oil<br>Marcol, oil<br>Shell Hydraulic oil<br>Isopropanol  | 371<br>98<br>2<br>2 | drums  L containers drums drums drums | Roane State Community<br>College via the Department<br>of General Services          |

## Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u> | Chemical   | Quan            | tity          |
|-------------|--|-----------------|---------------|
| 2/84        | Sulfuric acid  | 16              | kg            |
| 11/84       | o-Aminophenol-p-sulfonic acid Thioglycolic acid p-Anisidine acetyl o-Toluidine acetyl Dibromopropyl alcohol Cyanoacetanilide | 200<br>400<br>2 | g<br>kg<br>kg |
|             | Resorcinol monomethyl ether  | 75              | -             |
|             | Furil dioxime  | 20              | -             |
|             | Heptoyl chloride   | 300             | _             |
|             | 2-Hydroxy-1,4-Dimethyl benzene   | 250             |               |
|             | Methyl bromobenzoate   | 525             | -             |
|             | Methyl-p-aminophenol sulfate   | 200             | _             |
|             | Myristoyl chloride   | 100             | _             |
|             | P-Naphthol benzein   | 25              | _             |
|             | o-Nitrophenetole   | 100             |               |
|             | P-Nitrophenetole<br>P-Nitrophenylhydrazine   | 100<br>75       |               |
|             | P-Nitrophenylhydrazine   | 100             |               |
|             | hydrochloride  | 100             | 9             |
|             | P-Nitrosophenol sodium salt  | 125             | a             |
|             | Palmitoyl chloride   | 25              | _             |
|             | G-Phenylpropyl bromide   | 100             |               |
|             | Phloroglucinol   | 150             | _             |
|             | Propionamide   | 200             | g             |
|             | Tetra-N-Butylammonium iodide   | 400             | g             |
|             | Thio-B-Naphthol  | 50              | _             |
|             | Toluhydroguinone   | 250             | -             |
|             | P-Tolyl isocyanate   | 25              |               |
|             | Trimethylamine   | 250             | -             |
|             | Trimethylene bromide   | 200             | _             |
|             | Trimethylene chlorohydrin  | 25              |               |
|             | Xanthydrol<br>o-xylose   | 20<br>275       |               |
|             | 0-xy103e   | 2/5             | 9             |
| 5/85        | Lead carbonate   | 10              | ka            |
|             | Potassium bromide  | 25              |               |
|             | Sodium acetate   | 30              |               |
|             | Sodium peroxide  |                 | kg            |
|             | Tantalum pentoxide   |                 | kg            |
|             | Adipic acid  | 1               | kg            |
|             | o—Aminobenzenesulfonic acid  | 125             | _             |
|             | D-10-Camphorsulfonic acid  | 100             | _             |
|             | Chloroacetic acid  |                 | kg            |
|             | 3,5-Dinitrobenzoic acid  | 100             | _             |
|             | Iodoacetic acid  | 25              |               |
|             | B-Iodopropionic acid   | 25              | g             |


## Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u> | Chemical                           | Quantity |
|-------------|------------------------------------|----------|
| 5/85        | Naththalene-B-sulfonic acid        | 200 g    |
| cont.       | A-Naphthoic acid                   | 25 g     |
|             | o-Toluic acid                      | 800 g    |
|             | p-Toluic acid                      | 3 kg     |
|             | Undecylenic acid                   | 500 g    |
|             | Acenaphthene                       | 200 g    |
|             | Allyl formate                      | 100 g    |
|             | m-Aminophenol                      | 200 g    |
|             | o-Aminophenol                      | 100 g    |
|             | p-Aminophenol                      | 500 g    |
|             | p-Aminophenol hydrochloride        | 200 g    |
|             | Tert-Amylbenzene                   | 50 g     |
|             | n—Amyl bromide                     | 400 g    |
|             | o-Anisidine                        | 100 g    |
|             | p-Anisidine                        | 100 g    |
|             | Anthraquinone                      | 250 g    |
|             | Azoxybenzene                       | 100 g    |
|             | Benzil                             | 200 g    |
|             | Benzyl chloride                    | 2 kg     |
|             | A-Bromo-acetophenone               | 200 g    |
|             | Bromobenzene                       | 10 g     |
|             | Bromomesitylene                    | 200 g    |
|             | o-Bromophenol                      | 50 g     |
|             | 2-Bromopyridine                    | 100 g    |
|             | o-Bromotoluene                     | 50 mL    |
|             | Tert-butylbenzene                  | 25 g     |
|             | N-Butyl bromide                    | 250 g    |
|             | Isobutyl chloride                  | 200 g    |
|             | Isocaproyl chloride                | 50 g     |
|             | Cetyl bromide                      | 200 g    |
|             | Cetyl iodide                       | 100 g    |
|             | o-Chloroaniline                    | 500 g    |
|             | p—Chloroaniline                    | 100 g    |
|             | A-Chloromethylnaphthalene          | 500 g    |
|             | Crotonaldehyde                     | 500 g    |
|             | Cyanoacetamide                     | 200 g    |
|             | 1,2,5,6—Dibenzanthracene           | 1 g      |
|             | 9,10-Dibromoanthracene             | 50 g     |
|             | N-Dibutylamine                     | 5 kg     |
|             | Di-n-butyl Ketone                  | 25 g     |
|             | Dicyandiamide                      | 1 kg     |
|             | Diethyl Ketone                     | 3 kg     |
|             | Dimethylbenzylamine                | 200 g    |
|             | AS-Diphenylhydrazine hydrochloride | 25 g     |
|             | Di-p-tolyl Ketone                  | 25 g     |
|             | Petroleum ether                    | 8 L      |
|             | Ethyl anisate                      | 100 g    |
|             | Ethyl anthranilate                 | 100 g    |
|             | Ethyl B—Bromopropionate            | 100 g    |

## URAFT

## Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u>   | <u>Chemical</u>  | Quan | tity                |
|---------------|--|------|---------------------|
| 5/85<br>cont. | Ethylene chlorobromide<br>Ethylene chloride<br>Ethyl fumarate<br>Ethyl maleate | 1    | g<br>kg<br>kg<br>kg |
|               | Ethyl orthoformate   | 250  |                     |
|               | Lauryl bromide   | 700  | g                   |
|               | Methyl anthranilate  | 3    | kg                  |
|               | Methyl bromoacetate  | 400  | g                   |
|               | Methyl N-butylcarbinol   | 25   | g                   |
|               | Methyl N-butyrate  | 200  | g                   |
|               | o—Methylhydroxylamine hydrochloride  | 100  | g                   |
|               | Methyl malonate  | 500  | _                   |
|               | Methyl B-naphthyl Ketone   | 200  | -                   |
|               | Myristyl bromide   | 600  |                     |
|               | m-Nitroaniline   | 100  | _                   |
|               | o-Nitroaniline   | 200  |                     |
|               | 4-Nitro-1,2-Diaminobenzene   | 100  |                     |
|               | p-Nitrofluorobenzene   | 25   | -                   |
|               | Nitron   | 100  |                     |
|               | 1-Nitroso-2-napthol<br>o-Nitrotoluene  | 100  |                     |
|               | p-Nitrotoluene   | 250  | kg<br>~             |
|               | Phenanthraquinone  | 250  | _                   |
|               | Phenetole  | 500  |                     |
|               | Phenylbenzylcarbinol   | 25   | _                   |
|               | m-Phenylenediamine   | 250  |                     |
|               | DL-A-Phenylethylamine  | 100  |                     |
|               | DL-Phenylmethylcarbinol  |      | kg                  |
|               | Picoline   | 500  |                     |
|               | Propiophenone  | 1000 |                     |
|               | Iso-propylamine  |      | kg                  |
|               | Iso-propyl iodide  | 100  | g                   |
|               | Quinone  | 2    | kg                  |
|               | Trans-Stilbene   | 300  |                     |
|               | Strychnine sulfate   | 200  | g                   |
|               | Succinonitrile   | 200  |                     |
|               | Sucrose  | 150  |                     |
|               | m-Tolunitrile  | 25   |                     |
|               | o-Tolunitrile  | 25   |                     |
|               | p-Tolunitrile  | 100  |                     |
|               | N-Valeronitrile  | 100  |                     |
|               | Bromocresol green  |      | L                   |
|               | Buffer solution  | 64   | L                   |



| <u>Date</u> | Chemical  | Quantity   |
|-------------|---|--|
| 6/80        | 4-Iodo-n-xylene   | 250 g  |
| 6/80        | Silicic acid Sulfuric acid Ammonium thiocyanate Calcium chloride Sodium methylate Yttrium, metal 2-Amino-5-chlorobenzoic acid 1-Amino-2-naphthol-4-sulfonic acid o-Aminophenol-p-sulfonic acid B-Bromopropionic acid m-Chlorobenzoic acid Chlorosulfonic acid 3-Hydroxy-2-naphthoic acid 1-Malic acid DL-Mandelic acid Methoxyacetic acid o-Methoxybenzoic acid 3-Nitrosalicylic acid 5-Nitrosalicylic acid Salicylacetic acid Salicylacetic acid Salicylic acid Trimethylacetic acid Acetamide Allylthiourea N-Amyl chloride Tert-Amyl chloride Benzyl bromide Benzyl bromide Benzyl bromide N-Butyl-p-toluenesulfonate Chloroacetdiethylamide Chloroacetyl chloride Chloroacetyl chloride | 2 kg 3 kg 25 kg 30 g 25 g kg 25 g kg 30 g g kg 25 g kg 100 g g kg 100 g g g kg 100 g g g g g g 125 g g 100 g g g g g 125 g g 100 g g g g g 125 g g 100 g g g g g 100 g |
|             | A-Chloronaphthalene<br>2-Chloropyridine   | 3 kg<br>100 g  |
|             | Cinchonine, alkaloid o-Cresyl-p-toluenesulfonate Dicyclohexylamine Ethyl bromo-n-butyrate   | 50 g<br>200 g<br>1 kg<br>150 g   |
|             | Ethyl bromo-n-caproate Ethyl chloroacetate Ethyl naphthylamine m-Fluorochlorobenzene  | 75 g<br>250 g<br>100 g<br>25 g   |
|             | m-Fluorotoluene<br>o-Fluorotoluene  | 25 g<br>100 g  |

| <u>Date</u> | Chemical                                   | Quantity       |
|-------------|--|----------------|
| 8/81        | 2-Iodo-1,3-dimethylbenzene                 | 10 g           |
| cont        | Methyl adipate                             | 100 g          |
|             | Methyl bromopropionate                     | 300 g          |
|             | Methyl orthoacetate                        | 600 g          |
|             | 6—Methylquinoline                          | 25 g           |
|             | 8-Methylquinoline                          | 50 g           |
|             | Methyl toluate                             | < 1 kg         |
|             | Naphthyl acetate                           | 100 g          |
|             | p-Nitroacetanilide                         | 75 g           |
|             | A-Nitronaphthalene                         | 400 g          |
|             | m-Nitrophenol                              | 25 g           |
|             | Phenylethyl bromide                        | 300 g          |
|             | Phenylhydrazine                            | 500 g          |
|             | DL—Phenylmethylcarbinol                    | 9 kg           |
|             | Phenyl-p-tolyl ketone                      | 300 g          |
|             | Pinacol                                    | 500 g          |
|             | Propiophenone                              | 3 kg           |
|             | N-propyl carbonate                         | 100 g          |
|             | Propylene glycol                           | 8 L            |
|             | Thionaphthol                               | 225 g          |
|             | Ethyl bromopropionate                      | 100 g          |
|             | Glycine anhydride                          | 75 g           |
| 7/82        | Hydrofluosilicic acid                      | 3 kg           |
|             | Cadmium metal                              | 113 g          |
|             | Calcium chloride                           | 11 kg          |
|             | Lead sulfate                               | 1 kg           |
|             | Magnesium sulfate                          | 16 kg          |
|             | Potassium thiocyanate                      | 3 kg           |
|             | A-Bromo-n-valeric acid                     | 50 g           |
|             | o-Chlorobenzoic acid                       | 100 g          |
|             | Trans-Cinnamic acid                        | 300 g          |
|             | 2,4-Dinitrobenzene sulfonic acid           | 10 g           |
|             | Mucic acid                                 | < 1 kg         |
|             | Naphthalic anhydride acid<br>m—Toluic acid | 100 g          |
|             | Trichloroacetic acid                       | 2 kg           |
|             | Diacetone alcohol                          | 5 kg           |
|             | Myristyl alcohol                           | 250 g<br>50 g  |
|             | Phenylethyl alcohol                        | _              |
|             | Phenylpropyl alcohol                       | 1 kg<br>100 g  |
|             | 2-Aminopyrimidine                          | 300 g          |
|             | Iso—amyl bromide                           | 300 g<br>100 g |
|             | N-Amyl iodide                              | 300 g          |
|             | Benzalacetone                              | 300 g          |
|             | Benzidine dihydrochloride                  | 28 g           |
|             | Bromoacetyl bromide                        | 300 g          |
|             | Bromocyclohexane                           | 200 g          |
|             | N-Butyl bromide                            | 15 kg          |
|             |  |                |



| <u>Date</u>   | Chemical                                     | Quan       | <u>tity</u> |
|---------------|--|------------|-------------|
| 7/82<br>cont. | N—Butyl sebacate<br>Cellulose acetate        |            | kg          |
|               | Cellulose triacetate                         | 500        | _           |
|               | m—Chlorophenol<br>o—Chlorophenol             | 25         | g<br>kg     |
|               | Di-n-butylaniline                            | 300        | -           |
|               | p-Diethylaminobenzaldehyde                   | 25         |             |
|               | Diethyl-A-naphthylamine                      | 25         | _           |
|               | Ethylamine hydrochloride                     | 200        |             |
|               | Ethyl bromoacetate                           | 900        | _           |
|               | Ethylene bromide                             | 23         | _           |
|               | Ethylene chlorohydrin                        |            | kg          |
|               | Ethyl phosphate                              |            | kg          |
|               | Fluoranthene                                 | 200        |             |
|               | Fluorenone                                   | 100        | _           |
|               | N-Hexylphenylcarbinol                        | 40         | _           |
|               | o-Idobenzoyl chloride                        | 100        | _           |
|               | Methyl N-amyl Ketone                         |            | kg          |
|               | Methylethylamine hydrochloride               | 60         | g           |
|               | 4-Methyl-2-pentanol                          | 3          | kg          |
|               | Methyl phenylacetate                         | 250        | g           |
|               | Methyl—iso—thiocyanate                       | 50         | g           |
|               | 4-Methylumbelliferone                        | 25         | -           |
|               | 1,2-Naphthoquinone                           | 25         | -           |
|               | 1—Naphthyl isocyanate                        | 25         | -           |
|               | Pentaerythritol                              | 500        | _           |
|               | Phenanthrene                                 | 100        | _           |
|               | m-Phenetidine                                | 100        | _           |
|               | o-Phenetidine                                | 100        | _           |
|               | p-Phenetidine                                | 400        | -           |
|               | Phenyl acetate                               | 300        | _           |
|               | Phenyl—a—naphthylamine                       | 100<br>500 | _           |
| •             | Phthalyl chloride<br>p—Isopropylbenzaldehyde | 800        | _           |
|               | Isopropyl bromide                            | 500        |             |
|               | Propylene chloride                           | 750        | _           |
|               | N-Propyl sulfone                             | 25         | _           |
|               | p,p-Tetramethyldiaminodiphenylmethane        | 300        | _           |
|               | o-Tolidine dihydrochloride                   | 400        | _           |
|               | 1,2,3—Tribromopropane                        | 200        | _           |
|               | 1,1,1-Trichloroethane                        | 23         | Ĺ           |
|               | Xanthydrol                                   | 80         | g           |
| 8/82          | 1,1,1-Trichloroethane                        | 416        | L           |
|               | 2,2,4-Dihydroxybenzaldehyde                  | 700        |             |
| 4/83          | Ammonium acetate                             | 16         | kg          |
|               | Ammonium sulfide                             |            | L           |
|               | Antimony, metal                              | < 1        | kg          |



| <u>Date</u>   | <u>Chemical</u>   | Quantity   |
|---------------|---|--|
| 4/83<br>cont. | Calcium fluoride Ferrous sulphamate Lithium sulfate Mercurous nitrate Acetyl p-aminobenzoic acid B-Amino-1-naphthol-3,6-disulfonic acid DL-A-Aminophenylacetic acid 4-Amino-m-toluenesulfonic acid  | 5 kg<br>19 L<br>< 1 kg<br>< 1 kg<br>400 g<br>250 g<br>30 g<br>2 kg |
|               | p-Arsanilic acid DL-10-Camphorsulfonic acid Cyanuric acid A,B-Dibromosuccinic acid Diglycolic acid Furoic acid  | 200 g<br>200 g<br>300 g<br>75 g<br>200 g<br>200 g                  |
|               | Indole-3-acetic acid Acetyl thiourea B-Diethylaminoethyl alcohol 1-Aminoanthraquinone Benzoylacetone  | 2 g<br>200 g<br>1 kg<br>100 g<br>20 g                              |
|               | Benzyl thiocyanate Bromohydroquinone N-Butyl sulfone 2,4-Dibromophenol Di-isobutylene   | 600 g<br>50 g<br>50 g<br>100 g<br>4 kg                             |
|               | Dichloroacetyl chloride  1,3—Diphenylguanidine Ethyl,N—Butylmalonate Ethyl isothiocyanate Hexachloroethane Methyl thiocyanate   | 400 g<br>300 g<br>100 g<br>25 g<br>500 g<br>200 g                  |
|               | Phenyl isothiocyanate Propylene diamine Succinimide p,p—Tetramethyldiaminobenzophenone Xanthone Ion Exchange Resin  | 100 g<br>1 kg<br>300 g<br>25 g<br>50 g<br>3 kg                     |
| 4/84          | Molybdic acid Cesium oxide Potassium bifluoride Potassium, metal Potassium periodate Potassium thiocyanate Titanium oxide 1—Nitroanthraquinone—5—sulfonic acid Trichloroacetic acid Acetamide p—Aminophenol acetyl B—Dimethylaminoethyl alcohol | 1 kg 5 kg 5 kg < 1 kg 10 kg 5 g 800 g 1 kg 25 g 1 kg               |

# DRAET

| <u>Date</u> | Chemical                           | Quant | tity |
|-------------|------------------------------------|-------|------|
| 4/84        | p-Aminoacetophenone                | 50    | g    |
| cont.       | 2—Aminoanthraquinone               | 100   | g    |
|             | 3-Aminoquinoline                   | 50    | g    |
|             | 2-Aminothiazole                    | 1000  | _    |
|             | Benzenesulfonylamide               | 100   | g    |
|             | o-Bromophenetole                   | 125   | g    |
|             | p-Bromophenetole                   | 100   | g    |
|             | m—Chlorophenyl isocyanate          | 100   | g    |
|             | 2,4-Diaminophenol dihydrochloride  | 500   | g    |
|             | Dibenzalacetone                    | 25    | g    |
|             | 4,4—Dibromobiphenyl                | 200   | g    |
|             | 2-Diethylamino-1,4-dimethylbenzene | 50    | g    |
|             | 4,4-Dihydroxybenzophenone          | 50    | g    |
|             | 2,5-Dimethoxyaniline               | 75    | g    |
|             | Dimethyldihydroresorcinol          | 25    | g    |
|             | Phenyl cellosolve                  | 22    | kg   |
|             | Ethylene glycol, diacetate         | 8     | kg   |
|             | Hydantoin                          | 75    | g    |
|             | a-Hydrindone                       | 50    | g    |
|             | p-Hydroxybenzaldehyde              | 200   | g    |
|             | g-Hydroxybutronitrile              | 10    | _    |
|             | p-Hydroxyphenylglycine             | 100   | _    |
|             | o—Methoxybenzaldehyde              | 300   | _    |
|             | p—Phenylacetophenone               | 25    | _    |
|             | Isopropyl acetate                  | 100   | g    |
|             | Isopropyl acetate, purified        |       | L    |
|             | Tetramethylammonium chloride       | 1000  | g    |
|             | p—Toluamide                        | 75    |      |
|             | 1,2,3-Trichloropropane             | 200   | g    |
|             | Buffer solution                    | 23    |      |
|             | Ion Exchange resin                 | 8     | kg   |

DRAFT Copy



## OAK RIDGE NATIONAL LABORATORY

MARTIN MARIETTA

HISTORICAL CHEMICAL RELEASE REPORT
FOR

OAK RIDGE NATIONAL LABORATORY

DEPARTMENT OF ENVIRONMENTAL MANAGEMENT

ENVIRONMENTAL AND OCCUPATIONAL SAFETY DIVISON

OAK RIDGE NATIONAL LABORATORY

May 1986

NOTICE:

Internal Use Only

This document has not been given final patent clearance, and the dissemination of its information is only for official use. If this document is to be given public release, it must be reviewed in accordance with internal release procedures (D-8-5).

OPERATED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY

DRAFT

# FOR OAK RIDGE NATIONAL LABORATORY

DEPARTMENT OF ENVIRONMENTAL MANAGEMENT

ENVIRONMENTAL AND OCCUPATIONAL SAFETY DIVISON

OAK RIDGE NATIONAL LABORATORY

May 1986

## Historic Chemical Release Report for Oak Ridge National Laboratory

#### Introduction

This report provides information on the past use and disposition of toxic chemicals at the Oak Ridge National Laboratory (ORNL). As part of this effort, chemicals have been identified which were believed to have the greatest potential for environmental or public health impacts based on their toxicity and/or quantities used. Historically, the research activities at ORNL have utilized numerous chemicals in a wide-range of quantities, i.e., milligram to kilogram, milliliter to liter amounts. Approximately, 2,000 different chemicals were reviewed prior to formulating the final listing of 107 chemicals which are included in this report.

## Sources of Information

Several different information sources were considered in the preparation of this report. However, only those sources which were well documented have been utilized. These sources included the following:

- Annual Toxic Chemical Usage by ORNL Department Report 3063 (1980-1985)
- ORNL Hazardous Materials Usage Reconciliation Report 7010 (1980-1985)
- Purchase Requisitions with ORNL Account Numbers (1981-1985)
- Hazardous Waste Disposal Records (1982-1985)
- Asbestos Containing Waste Materials Disposal Records
- Waste Management Operating Records for ORNL Waste Treatment Systems
- Plant & Equipment Operating Records for Grounds Maintenance
- PCB Tracking System

Because of the limitation to use only documented records, in most cases, it was not possible to provide a good retrospective database of more than five years.

## <u>Historical Chemical Findings</u>

Information for 107 chemicals is provided in the Attachments and Appendices. Attachment I is comprised of the completed Chemical Release Data Forms on each chemical. Attachment II provides available information on the estimated chemical usage for each of the 107 items. Finally, Attachment III provides a summation of ORNL surplus chemicals and other related activities. The Appendices serve to provide some additional information on specific chemical releases of significance.

# ATTACHMENT I Chemical Release Data Forms

| Chemical | Name: | Acetone |
|----------|-------|---------|
|          |       | claaner |

Uses: Solvent, cleaner, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constituent present in spill material that resulted in a fish-kill in White Oak Creek in 1983. See Appendix I for more details.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department -Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical    | Name: | Aceton | itr | i | le |
|-------------|-------|--------|-----|---|----|
| • • • • • • |       |        | _   | _ |    |

Uses: Solvent, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name: Acrylamide<br>Uses: Ion exchange columns,<br>Solid X Liquid<br>Listed as Toxic: Yes X | chemical laboratory uses Gas No |
|--|---------------------------------|
| Status of Environmental Comp   | liance (Past and Present):      |

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Nam | e: A  | luminu | m nit | rate |          |    |
|---------|-------|-------|--------|-------|------|----------|----|
| Uses:   | Chemi | cal 1 | aborat | ory u | ises |          |    |
| Solid   | X     | _ Liq | uid    |       | Gas  |          |    |
| Listed  | as To | xic:  | Yes    |       | No _ | <u> </u> | _• |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Ammonia, anhydrous
Uses: Refrigerant, chemical reagent, blueprinting
Solid Liquid Gas X
Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Ammonium hydroxide
Uses: Cleaners, chemical laboratory uses
Solid Liquid X Gas \_\_\_\_\_.
Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Na | ıme: A | mmoniu | id mu     | fluori | de |
|---------|------|--------|--------|-----------|--------|----|
| Uses:   | Elec | tropla | ting ( | opera     |        |    |
| Solid   | X    | Liq    | uid _  |           | Gas .  |    |
| Listed  | as   | Toxic: | Yes    | <u> X</u> | _ No _ | ·  |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

|          |        | : Ammoniu | m nitrate |     |
|----------|--------|-----------|-----------|-----|
| Uses:    |        |           | _         |     |
|          |        | _Liquid   | Gas       |     |
| listed [ | as Tox | ic: Yes   | No        | Χ . |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Asbestos - Containing materials
Uses: Insulation
Solid X Liquid Gas Listed as Toxic: Yes X No Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Asbestos has been used at ORNL and there are various on-going asbestos removal projects. The study conducted (Aug. 13 through Sept. 21, 1984) for Non-radiological Process Wastewater Characterization indicated a minimum of <0.3 and a maximum of 3.00 (with a mean value of 0.35) Million Fibers per Liter (MFL). The flume location which served as a reference for this study indicated <0.3 MFL. No environmental impact is presently known or suspected due to asbestos.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Asbestos-containing waste disposal records.

| Chemica | ll Nam | e: B  | larium | octa | ahydrate | 7 |   |  |
|---------|--------|-------|--------|------|----------|---|---|--|
| Uses:   | Chemi  | cal 1 | abora  | tory | uses     |   |   |  |
| Solid   | Χ      | Lia   | uid    |      | Gas _    |   |   |  |
| Listed  | as To  | xic:  | Yes    |      | No _     | X | • |  |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

|        |        |         | hyl carbamat | e (Baygon) |
|--------|--------|---------|--------------|------------|
| Uses:  | Insect | icide   |              |            |
| Solid  | X      | Liquid  | Gas          | ·          |
| listed | as Tox | ic: Yes | No _         | X•         |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemica | 11 1 | lame:  | Benze | ne  |       |       |      |
|---------|------|--------|-------|-----|-------|-------|------|
| Uses:   | So   | lvent, | chemi | cal | labor | atory | uses |
| Solid   |      |        | iquid |     | G     | as    |      |
| Listed  | as   | Toxic  | : Yes | X   | N N   | 0     | •    |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constituent present in spill material that resulted in a fish-kill in White Oak Creek in 1983. See Appendix I for more details.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica |     |      |              |       |      |        |          |   |
|---------|-----|------|--------------|-------|------|--------|----------|---|
| Uses:   | Che | emic | <b>a</b> 1 1 | abora | tory | uses   |          |   |
| Solid   | X   | ζ    | Lic          | uid _ |      | Gas    |          | _ |
| Listed  | as  | Tox  | ic:          | Yes   | X    | _ No _ | <u> </u> |   |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | il N | lame: | С    | alcium | hypo | ochlorit | :e   |   |
|---------|------|-------|------|--------|------|----------|------|---|
| lises:  | B16  | each, | , ch | emical | 1 ab | oratory  | uses |   |
| hilo2   | >    | (     | Lia  | uid    |      | _ Gas    |      | _ |
| Listed  | as   | Tox   | c:   | Yes    | Χ    | No       |      |   |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Carbon monoxide
Uses: Ecological research uses, synthesis
Solid Liquid Gas X
Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Carbon tetrachloride

Uses: Cleaner, solvent, chemical laboratory uses

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been one carbon tetrachloride spill 3.78 L (1 gal) in 1984. However, it occurred indoors and there was no evidence of environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

|           | ame: Chlord | ane |         |   |
|-----------|-------------|-----|---------|---|
| Uses: Ins | ecticide    |     |         |   |
| J U       | Liquid      | X   | _ Gas _ |   |
| listed as | Toxic: Yes  | X   | No      | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemica | ] | Name: | Ch | lori | ine, | gas | ; |
|---------|---|-------|----|------|------|-----|---|
| Heac.   |   |       |    |      |      |     |   |

Uses: Treatment of Sewage Treatment Plant effluent

Solid Liquid Gas X Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

An unknown amount of chlorine and helium were released at Bldg. 4500S on 2/26/85. There were no known environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department -Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- ORNL Waste Management operating records.

|       | 1 Name:  |         |       |         |        |
|-------|----------|---------|-------|---------|--------|
| Uses: | Solvent, | chemica | al la | borator | y uses |
| Solid |          | iquid   | X     | Gas     | •      |
|       | ac Tovic | · Vac   | Y     | _ No _  |        |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al Name: | Chlor | pyrifos | (Bolt | ) |
|---------|----------|-------|---------|-------|---|
| Uses:   | Insectic | ide   |         |       |   |
| Solid   | L        | iquid | X       | _ Gas |   |
|         | as Toxic | : Yes | Χ       | No    | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Chromic acid

Uses: Cleaner, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No .....

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Coal oil mixture

Uses: R&D work in coal conversion programs

Solid Liquid X Gas Listed as Toxic: Yes No X

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Coal Tar Pitch

Uses: Roofing material, R&D coal conversion programs

Solid X Liquid X Gas Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing; (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Cyclohexane

Uses: Solvent, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No ................

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Nar |          | on |     |   |
|--------------|----------|----|-----|---|
| Uses: Inse   | cticide  |    | •   |   |
|              | Liquid   | Χ  | Gas |   |
| Lictor as To | vice Yes | X  | No  | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: 2,4 - Dichloropenaxy acetic acid (2,4-D)

Uses: Herbicide

Solid Liquid X Gas Listed as Toxic: Yes X No .....

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Diethylbenzene

Uses: Solvent, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Uses: Chemical synthesis, chemical laboratory uses Solid X Liquid Gas Listed as Toxic: Yes No X | Chemical Name:  | Diethylenetr | iamine  | pentacetic | sodium | salt |
|---|-----------------|--------------|---------|------------|--------|------|
| Solid X Liquid Gas No X   | Uses: Chemical  | synthesis, c | hemical | laboratory | uses   |      |
| licted as Toxic: Yes NO X   | Solid X L       | iquid        | Gas     |            |        |      |
| Listed as locate: les no  | Listed as Toxic | : Yes        | No      | <u>X</u>   |        |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemical Name: 1<br>Uses: Solvent, 1 | iquid scintil      | lation counting, | chemical | 1aboratory | uses |
|--------------------------------------|--------------------|------------------|----------|------------|------|
| Solid Liq<br>Listed as Toxic:        | uid X G<br>Yes X N | as<br>0          |          |            |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name | e: D | isulfo | oton | (Disyston | G) |  |
|---------|--------|------|--------|------|-----------|----|--|
| Uses:   |        |      |        |      |           |    |  |
| Solid_  | X      |      |        |      | _ Gas     |    |  |
| listed  | as Tox | (ic: | Yes    | Χ    | No        | •  |  |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Dodecane

Uses: Solvent

Solid Liquid X Gas Listed as Toxic: Yes No X

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al Name: | Sodium s   | alt organo    | o-phosphor | ic acid, | Merceptobenzotriazolo |
|---------|----------|------------|---------------|------------|----------|-----------------------|
|         |          | (Fndcor)   |               |            |          |                       |
| Uses:   | Corrosi  | ve inhibit | or used in    | n cooling  | towers   |                       |
| Solid   |          | Liquid     | <u>X Gas </u> |            |          |                       |
| Listed  | as Toxi  | c: Yes     | No _          | X          |          |                       |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Results of a toxicity test using cooling tower effluents have indicated detrimental effects on acquatic test organisms. The use of this chemical as a corrosive inhibitor in the cooling towers has been terminated.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Epoxy resins

Uses: Coatings, adhesives, potting of electronic circuits

Solid Liquid X Gas Listed as Toxic: Yes No X

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Ethyl acetate

Uses: Solvent, paint, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No .

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name:<br>Uses: Solvent, | Ethyl alcoleaner/ | cohol<br>sterilizer, | chemical | and | biological | laboratory | uses |
|----------------------------------|-------------------|----------------------|----------|-----|------------|------------|------|
| Solid Li<br>Listed as Toxic:     |                   |                      |          |     |            |            |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constitunet present in spill material that resulted in a fish-kill in White Oak Creek in 1983. See Appendix I for more details.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name:  | Ethyl, alcoh   | nol, denat | ured       |            |      |
|-----------------|----------------|------------|------------|------------|------|
| Uses: Solvent,  | cleaner, che   | emical and | biological | laboratory | uses |
| Solid Li        | iquid <u>X</u> | _ Gas      | •          |            |      |
| Listed as Toxic | : Yes <u>X</u> | No         | <b>_•</b>  |            |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | l Name:<br>Solvent, | Ethyl anest | ether<br>hetic, | chemical | laboratory    | uses |
|---------|---------------------|-------------|-----------------|----------|---------------|------|
| Solid   | as Toxic            | iauid       | Χ               | Gas      | <del></del> . |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Ethylene glycol

Uses: Antifreeze, coolant, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes No X .

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Four ethylene glycol spills took place due to mechanical failures or operator error on 2/13/80, 10/26/81, 7/2/84, and 10/24/85. These spills ranged in quantities from 7.57 L (2 gal) to 75,700 L (20,000 gal). There was no evidence of environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Ethylene oxide

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | ll Name | e: F  | erric | chlo | oride |      |
|---------|---------|-------|-------|------|-------|------|
| Uses:   | Chemi   | cal 1 | abora | tory | uses  |      |
| Solid   | X       | _ Liq | uid _ |      | Gas   | <br> |
| listed  | as To   | χic:  | Yes   | Χ    | No    | •    |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica |      |         |       |      |       |       |
|---------|------|---------|-------|------|-------|-------|
| Uses:   | Chem | nical 1 | abora | tory | uses  |       |
|         |      | Liq     |       |      | _ Gas |       |
| Listed  | as T | oxic:   | Yes   | • Х  | — No  | <br>• |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Formaldehyde
Uses: Sterilant, tissue fixative, chemical laboratory uses
Solid \_\_\_\_\_Liquid \_\_X \_\_Gas \_\_\_\_.
Listed as Toxic: Yes \_\_X \_\_ No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica |    |        |         |       |        |          |  |
|---------|----|--------|---------|-------|--------|----------|--|
| Uses:   | Ch | emical | labora  | atory | uses   |          |  |
| Solid   |    | L.     | iquid _ | X     | Gas    |          |  |
| Listed  | as | Toxic  | : Yes   |       | _ No _ | <u> </u> |  |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Hexane

Uses: Solvent, paints, chemical laboratory uses

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

On 4/17/84, 0.47 L (1 pint) of hexane was spilled at Bldg. 7001. There was no environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical | Name:   | Hydrof | luorio | : acid: |   |
|----------|---------|--------|--------|---------|---|
| Solid    |         | iquid  |        | _ Gas _ |   |
| Listed a | s Toxic | : Yes  | X      | No _    | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Hydrogen peroxide

Uses: Bleach, disinfectant, chemical uses

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica |     |        |       |       |      |  |
|---------|-----|--------|-------|-------|------|--|
| Uses:   | Che | emical | labor | atory | uses |  |
| Solid   |     | Li     | quid  | X     | Gas  |  |
| Listed  | as  | Toxic  | Yes   | X     | — No |  |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name:  | Isopropy | /l ald | coho 1    |            |            |      |
|-----------------|----------|--------|-----------|------------|------------|------|
| Uses: Solvent,  | paints,  | disi   | nfectant, | chemical   | laboratory | uses |
| Solid Li        | quid     | Χ      | Gas       | ·          |            |      |
| Listed as Toxic | Yes      | X      | No        | <b>_</b> • |            |      |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name:                |              |           | richloromethyl |
|-------------------------------|--------------|-----------|----------------|
| •                             | benzhydrol ( | Kelthane) |                |
| Uses: Insectici<br>Solid X Li |              | Gas       | •              |
| Listed as Toxic:              | Yes X        | No        | -              |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant and Equipment operating records.

| Chemical | Name: | Lacquer | thinner |
|----------|-------|---------|---------|
| 5        |       | +       |         |

Uses: Paints, coatings

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilalation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.

٠: ٠٠

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant and Equipment operating records.

| Chemica | al N | ame | : Le | ead | she  | et  |      |        |          |     |
|---------|------|-----|------|-----|------|-----|------|--------|----------|-----|
| Uses:   | Sol  | der | ing, | rac | diat | ion | shie | lding, | all      | oys |
| hilo2   | X    |     | lia  | biı |      |     | Gas  |        | <u>•</u> |     |
| Listed  | as   | Tox | ic:  | Yes | 5    | Χ   | No   |        | •        |     |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

During the operation of SWSA 6, mixed wastes containing lead were disposed by land burial. The implication of this information is that portions of SWSA 6 should be classified as RCRA disposal units.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

COLOR DE CARROL EN COMPANY DE CONTRACTOR DE

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant and Equipment operating records.

| Chemica | ıl N | ame:  | Mag  | <sub>j</sub> nes i | um n | itrate  |      |
|---------|------|-------|------|--------------------|------|---------|------|
| Uses:   | Che  | mical | 1 at | orat               | cory | uses    |      |
| Solid   | Χ    | L     | iqui | d                  |      | _ Gas _ | <br> |
| Listed  | as   | Toxic | : Y  | es_                | Χ    | No      | •    |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical | Name: | Malathior |
|----------|-------|-----------|
|----------|-------|-----------|

Uses: Insecticide

Solid Liquid X Gas Listed as Toxic: Yes X No . .

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant and Equipment operating records.

| Chemical Name: Mercury       | •   |    |
|------------------------------|---|----|
| Uses: Lamps, batteries, gaug | ges, instruments, chemical laboratory use | 32 |
| Solid Liquid X (             | _   |    |
| Listed as Toxic: Yes X       | No•                                       |    |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There were twelve cases of recorded mercury spills at ORNL ranging from trace amounts to 0.1 kg (0.22 lb) between 1980 and 1985. There were no known environmental impacts from these spills.

There have been occasions at ORNL when pockets of mercury were discovered at pipes or under tiles. These small quantities are attributed to the processes which were undertaken at ORNL to support the Y-12 thermonuclear weapons program. These operations took place in the 1950s and 1960s at Bldgs. 4501, 3503, and 3592. Though there is not an accurate measure of the mercury loss at ORNL, operating personnel have estimated losses of 907 - 1361 kg (2000-3000 1b) due to spills and leakage.

In a recent study, soil samples collected around Bldg. 4501 contained mercury concentrations ranging from 0.05 to 4.4 ppm. However, one sample was as high as 465 ppm. Soil samples near Bldg. 3503 contained mercury concentrations ranging from 0.8 to 25 ppm and near Bldg. 3592 the samples ranged from 4.1 to 320 ppm.

Chemical Name: Mercury

Page 2

Known or Suspected Environmental Impacts: (cont.)

Though there were individual fish samples with higher concentrations, all average values of mercury concentrations in Clinch River fish were below the Food and Drug Administration action level of 100 ng/g.

Mercury concentrations in water collected at White Oak Dam, White Oak Creek, and Melton Hill Dam were all higher than the Tennessee stream criteria. (Source: Environmental Monitoring Report, ORNL-6209).

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Methyl alcohol

Uses: Solvent, paint, fuel, coolant, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical Name:  | Methyl e | ethyl | keto  | ne       |            |      |
|-----------------|----------|-------|-------|----------|------------|------|
| Uses: Solvent,  | paints,  | coat  | ings, | chemical | laboratory | uses |
| Solid Li        | iquid    | Χ     | Gas   | •        |            |      |
| Listed as Toxic | : Yes    | X     | No    | ·        |            |      |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts. \_ \_ \_

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: p,p' - Diaminodiphenylmethane Uses: Plastic coatings

Solid X · Liquid Gas Listed as Toxic: Yes No X ·

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name:  | Methyle | ne ch' | lorid | e          |          |            |      |
|---------|----------|---------|--------|-------|------------|----------|------------|------|
| Uses:   | Solvent, | paints, | coat   | ings, | degreaser, | chemical | laboratory | uses |
| Solid   | Li       | iquid   | Χ      | Gas   | •          |          |            |      |
| Listed  | as Toxic | : Yes _ | X      | No _  | •          |          |            |      |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constituent present in spill material that resulted in a fish kill in White Oak Creek in 1983. See Appendix I for more details.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Metex Stripper
Uses: Metal cleaning
Solid Liquid X Gas
Listed as Toxic: Yes No X .

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Micro-bio-Treat [n-alkyl dimethyl benzylammonium

(Bis (Tri-n-butyltin) oxide]

Uses: Control algae growth in cooling towers

Solid Liquid X Gas Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Results of toxicity test using cooling tower effluents have indicated detrimental effects on acquatic test organisms. The use of this chemical for controlling algae growth in cooling towers has been terminated.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant and Equipment operating records.

| Chemica | 1 Name:  | Naptha  |      |       |          | 2 - b b    |      |
|---------|----------|---------|------|-------|----------|------------|------|
| Uses:   | Solvent, | paints, | coat | ings, | chemical | laboratory | นรยร |
| Solid   | Li       | iquid   | Χ    | Gas   | •        |            |      |
| Listeɗ  | as Toxic | : Yes   | X    | No _  | ·        |            |      |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Nickel Chloride
Uses: Metal plating, chemical laboratory uses

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name | e: Ni  | ckel  | sulfat    | :e   |            |      |
|---------|--------|--------|-------|-----------|------|------------|------|
| Uses:   | Metal  | plat.  | ing,  | chemica   | al 1 | laboratory | uses |
| Solid _ | X      | _ Liqu | _ biı |           | Gas  | ·          |      |
| Listed  | as Tox | (ic:   | Yes   | <u> X</u> | No   | <u> </u>   |      |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Nitric acid

Uses: Cleaning, metal prep., chemical reagent

Solid Liquid X Gas Listed as Toxic: Yes X No \_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Five different nitric acid spills, ranging from trace amounts to 3.79 L (1 gal), took place indoors. There was no environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al N | ame:   | Nitric | oxid  | le   |  |   |
|---------|------|--------|--------|-------|------|--|---|
| Uses:   | Che  | mical  | labora | tory  | uses |  | • |
| Solid   |      | Li     | quid   | Gas _ | X    |  |   |
| Listed  | as   | Toxic: | Yes    | X     | No   |  | - |

Status of Environmental Compliance (Past and Present):

At present all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs and trichlorethylene and these are monitoring only requirements. The primary reason for this is that existing data do not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica                        | al Name: | : Acetylph | osphoramid | othioic | acid, | 0,5-dimethyl | ester | (Orthene |
|--------------------------------|----------|------------|------------|---------|-------|--------------|-------|----------|
| Uses:                          | Insecti  | cide       |            |         |       |              |       |          |
| Solid                          | Χ        | Liquid     | Gas _      | •       |       |              |       |          |
| ${	t Listed}^{oldsymbol{	au}}$ | as Toxi  | c: Yes     | No         | _X•     |       |              |       |          |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemica | al Na | ame: P  | araro | sanil | ine   |      |    |
|---------|-------|---------|-------|-------|-------|------|----|
| Uses:   | Dye   | s, chem | ical  | labor | atory | uses |    |
| Solid   | _     | Liq     | uid _ | Χ     | Gas   |      | ·  |
| Listed  | as    | Toxic:  | Yes   |       | No    | X    | _• |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.

9 18 0 4 4 5 TRES 1

| Chemica |    |        |        |        |         |         |
|---------|----|--------|--------|--------|---------|---------|
| Uses:   | So | lvent, | chemic | cal la | aborato | ry uses |
| Solid   |    | Li     | quid   | X      | Gas _   |         |
| Listed  | as | Toxic  | : Yes  | X      | No _    |         |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

· Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name:  | Perchlo | oric | acid |  |
|---------|----------|---------|------|------|--|
| Uses:   | Chemical | labora  | tory | uses |  |
| Solid   | Li       | iquid   | X    | Gas  |  |
| Listed  | as Toxic | : Yes   | X    | No   |  |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.

....

Chemical Name: Petroleum ether
Uses: Solvent, chemical laboratory uses
Solid Liquid X Gas
Listed as Toxic: Yes X No

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al N | Name: P  | heno l  |          | •       |        |      | •      |     |
|---------|------|----------|---------|----------|---------|--------|------|--------|-----|
| Uses:   | Di   | sinfecta | ınt, ch | emica    | l labor | ratory | uses | 3      |     |
| Solid _ | Х    | Liq      | uid     |          | Gas _   | •      |      |        |     |
| Listed  | as   | Toxic:   | Yes _   | <u>X</u> | No      | ·      |      |        |     |
| Status  | of   | Environ  | mental  | Comp     | iance   | (Past  | and  | Presen | t): |

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemic  | al Name | e: Ph  | nenolic  | resin      |               |          |         |       |
|---------|---------|--------|----------|------------|---------------|----------|---------|-------|
| Uses:   | Parts   | fabr   | ication, | potting    | for e         | lectro   | nic cir | cuits |
| Solid . | Х       | _ Liqu | uid      | Gas _      | . <del></del> | <b>•</b> |         |       |
| Listed  | as To   | xic:   | Yes      | No         | <u>X</u> .    | •        |         |       |
| Status  | of En   | viron  | mental C | Compliance | (Pas          | t and    | Present | ):    |

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al Name | e: Pi | nosph | noric  | acid   |        |           |     |
|---------|---------|-------|-------|--------|--------|--------|-----------|-----|
| Uses:   | Metal   | prep  | ., c  | leanir | ig, ch | emical | reage     | ent |
| Solid   |         | Lia   | uid   | X      |        |        | <b></b> • |     |
| Listed  | as To:  | xic:  | Yes   | X      | _ No   |        | •         |     |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

|                 |    |     |      |     |   |       | biphenyls | (PCBs |
|-----------------|----|-----|------|-----|---|-------|-----------|-------|
|                 |    |     |      |     |   |       | quipment  |       |
| Solid<br>Listed |    |     | Liqu | id  | X | _ Gas | •         |       |
| Listed          | as | Tox | ic:  | Yes | X | No    | •         |       |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

On May 6, 1981, and April 4, 1984, rupturing transformer incidents took place. In each case, oil from transformers were sprayed on desk tops, floors, and nearby personnel. Incidents occurred inside the buildings. Samples were taken and areas were decontaminated. In addition to these, two PCB oil spills and three light ballast leakages containing PCBs have occurred. There have been no known environmental consequences from these incidents. Further information on PCB related incidents is included in Appendix II.

According to the Environmental Monitoring Report, ORNL-6209 for 1984, PCB concentration in Clinch River fish at CRM 20.8 on the average were very close to background level and were below the 2  $\mu$ g/g wet weight tolerance level designated by the Food and Drug Administration. The total PCB concentration in water collected about Melton Hill Dam was 0.00014 ppm.

Chemical Name: Polychlorinated biphenyls (PCBs)

Page 2

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.

- PCB Tracking System.

| Chemica | al Name | e: Po | tass | ium cy | anid | le         |      |
|---------|---------|-------|------|--------|------|------------|------|
| Uses:   | Metal   | plati | ng,  | chemic | al 1 | laboratory | uses |
| Solid   | X       | Liau  | id   |        | Gas  |            |      |
| Listed  | as To   | Ric:  | Yes  | X      | No   |            |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Potassium hydroxide

Uses: Plating, cleaning, chemical reagent

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

On May 25, 1984, and April 9, 1986, 37.85 L (10 gal) and 1.9 L (0.5 gal) of potassium hydroxide were spilled within buildings 7000 and 7920. There were no environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- ORNL Waste Management operating records.

| Chemica | al Name: | Pramit   | :01 |       |       |
|---------|----------|----------|-----|-------|-------|
| Uses:   | Herbici  | de       |     |       |       |
| Solid _ | Χ        | Liquid _ |     | Gas . | <br>' |
| isted   | as Toxi  | c: Yes   |     | No    | •     |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant and Equipment operating records.

|        | al Name: |        |       |          |      |
|--------|----------|--------|-------|----------|------|
| Uses:  | Solvent, | chemic | al la | boratory | uses |
| Solid  | L        | iquid  | X     | Gas      | ·    |
| Listed | as Toxic | : Yes  | X     | No       | •    |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al Name | : Ronstar | •   |    |   |
|---------|---------|-----------|-----|----|---|
| Uses:   | Herbic  | ide       |     |    |   |
| Solid   | X       | Liquid    | Gas |    |   |
| listed  | as Tox  | ic: Yes   | No  | _X | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemica         | 1 Name | : N-Phosp                  | honome | ethyl | glycine       | isopropylamine | salt | (Round-up) |
|-----------------|--------|----------------------------|--------|-------|---------------|----------------|------|------------|
| Uses:<br>Solid_ | Herbic | ide<br>_Liquid<br>ic: Yes_ | X      | Gas _ | •             |                |      |            |
| Listed          | as lox | ic: Yes _                  |        | NO _  | <del></del> • |                |      |            |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Silicon tetrafluoride
Uses: Chemical laboratory uses
Solid Liquid Gas X
Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | al Name | : Silvex |      | • |
|---------|---------|----------|------|---|
| Uses:   | Herbic  | ide .    |      |   |
| Solid   | Χ _     | Liquid _ | Gas  |   |
| listed  | as Tox  | ic: Yes  | X No | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemica | 17 1 | lame:  | Sodi   | um bif | luoride | • |
|---------|------|--------|--------|--------|---------|---|
| Uses:   | Che  | emical | llabo  | ratory | uses    |   |
| Solid   | >    | ( L    | .iquid |        | Gas     |   |
| Listed  | as   | Toxic  | ∷ Ye   | s X    | No _    | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) properly protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name | e: Sodium | n cyanide | 1 - houstony |      |
|---------|--------|-----------|-----------|--------------|------|
| Uses:   | Metal  | plating,  | cnemical  | laboratory   | uses |
| Solid_  | X      | _Liquid _ | G         |              |      |
| Listed  | as To  | kic: Yes  | X No      | o•           |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Lifelifeat Maile. Soutain dictionis. | Chemical | Name: | Sodium | dithionite |
|--------------------------------------|----------|-------|--------|------------|
|--------------------------------------|----------|-------|--------|------------|

Uses: Silver recovery, chemical laboratory uses

Solid X Liquid Gas Listed as Toxic: Yes X No .......

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

On 7/24/86 a sodium dithionite related spontaneous chemical reaction occurred at Bldg. 4500S. The vessel containing the reaction was removed from the building to an environmentally safe location and the incident resolved. No environmental impact is known from this accident.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Silver Recovery processing records.

| Chemica | al N | lame | : S | odium | hex | ametaph | osph | ate  |
|---------|------|------|-----|-------|-----|---------|------|------|
| Uses:   | Cle  | eani | ng, | chemi | cal | laborat | ory  | uses |
| Solid   | X    | (    | Liq | uid   |     | Gas     |      |      |
| Listed  | as   | Tox  | ic: | Yes   |     | No _    | X    |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name | 2: So | dium  | Hydr | oxide    |         |
|---------|--------|-------|-------|------|----------|---------|
| Uses:   | Clean  | ing,  | plati | ng,  | chemical | reagent |
| Solid   | Х      |       |       |      | Gas      | ·       |
| listed  | as Tox | cic:  | Yes_  | X    | No       | •       |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

Sodium hydroxide spilled during the filling of a tank inside Bldg. 3004 via a line from a tanker outside. Even though the tank was diked, 3.78 L (1 gal) of material was spilled outside the containment dike. Spill and washings caused NPDES noncompliance; however, it was only a temporary environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Réport 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal operations.
- Waste Management operating records.

| Chemica | 1 Nam | e: S | odium | nitr | ate . |   |
|---------|-------|------|-------|------|-------|---|
| Uses:   |       |      |       | tory |       |   |
|         | X     |      |       |      | _ Gas |   |
| listed  | as To | xic: | Yes   | Χ_   | No_   | · |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica |     |       |       |     |      |      |        |
|---------|-----|-------|-------|-----|------|------|--------|
| Uses:   | Che | emica | l lab | ora | tory | uses |        |
| Solid   | X   | L     | iqui. | d   |      | Gas  | <br>   |
| Listed  | as  | Toxid | :: Y  | es  | Χ    | No   | <br>_• |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Subdue E2

Uses: Fungicide

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Sulfur hexafluoride

Uses: dielectric, tracer gas

Solid Liquid Gas X Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemical | Name: | Sulfur | ic | acid |
|----------|-------|--------|----|------|
|          |       |        |    |      |

Uses: Cleaning, chemical laboratory uses
Solid Liquid X Gas
Listed as Toxic: Yes X No

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

On March 24, 1980, a 60.56 L (16 gal) spill of sulfuric acid was traced to Bldg. 3544. This spill caused an NPDES permit violation for pH. However, this was a temporary environmental impact.

Indoor spills on March 19, 1981, March 30, 1980, and April 11, 1986, of 22.71 L (6 gal); 3.79 L (1 gal); and 37.85 L (5-10 gal), respectively, of sulfuric acid were reported; however, there was no cause to believe they created an environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Waste Management operating records.

Chemical Name: 4'-Bromo (1,1'-biphenyl)-4-yl (Talon)
Uses: Rodenticide
Solid X Liquid Gas No X.
Listed as Toxic: Yes No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

- Hazardous waste disposal records.
- Plant & Equipment operating records.

Chemical Name: Tetrachloroethylene

Uses: Cleaner, degreaser, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No .

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Tetrahydrofuran
Uses: Solvent, chemical laboratory uses
Solid Liquid X Gas
Listed as Toxic: Yes X No \_\_\_\_\_.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride; tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Uses: | al Name: Toluene<br>Solvent, paints,<br>scintillation cou |       | chemical | laboratory | uses, | liquid, |
|-------|---|-------|----------|------------|-------|---------|
| Solid | Liquid<br>as Toxic: Yes                                   | X Gas | ·•       |            |       |         |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

During the operation of SWSA 6, mixed wastes containing toluene were disposed by land burial. The implication of this information is that portions of SWSA 6 should be classified as RCRA disposal units.

Known or Suspected Public Health Impacts:

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constituent present in spill material that resulted in a fish-kill in White Oak Creek in 1983. See Appendix I for more details. On May 11, 1983, 5.68 L (1.5 gal) of toluene was spilled at Bldg. 4500N. No environmental impact resulted.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include one or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage by ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Tributyl phosphate
Uses: Solvent, chemical laboratory uses
Solid Liquid Gas
Listed as Toxic: Yes No X

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

A trace amount of this chemical was spilled at Bldg. 7001 on 11/28/84. There was no environmental impact.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Uses: | al Name:<br>Solvent,       | paints, | coatings | e<br>, chemical | 1aboratory | uses, | liquid |
|-------|----------------------------|---------|----------|-----------------|------------|-------|--------|
| bilo2 | scintilla<br>L<br>as Toxic | iauid   | X Gas    | ·               |            |       |        |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

- No known or suspected environmental impacts.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | A F | lame:  | Trichl | oror | nethyls | ilan    | ıe   |
|---------|-----|--------|--------|------|---------|---------|------|
| Uses:   | So  | lvent, | chemic | cal  | laborat | ory     | uses |
| Solid   |     | L      | iquid  | X    | Gas     |         |      |
| Listed  | as  | Toxic  | : Yes  |      | No      | $-\chi$ | •    |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Triethanolamine

Uses: Cleaner, degreaser, chemical laboratory uses

Solid Liquid X Gas Listed as Toxic: Yes X No ......

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

Chemical Name: Urethane

Uses: Solvent, chemical laboratory uses

Solid X Liquid Gas No X.

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

|        | 1 Name:  |         |    |     |   |
|--------|----------|---------|----|-----|---|
| Uses:  | Solvent, | cleani  | ng |     |   |
| Solid  | Li       | iquid _ | Χ  | Gas |   |
| Listed | as Toxic | : Yes   | X  | No  | • |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| Chemica | 1 Name | : Vydat | e   |    |
|---------|--------|---------|-----|----|
| Uses:   | Insect | icide   | _   |    |
|         |        | Liquid  | Gas |    |
| listed  | as Tox | ic: Yes | No  | Χ. |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

There has been no known or suspected environmental impact due to the historic releases of this chemical.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

- Hazardous waste disposal records.
- Plant & Equipment operating records.

| Chemical Name:<br>Uses: Cleaner, | Xylene<br>degreaser, | chemical  | laboratory | uses |
|----------------------------------|----------------------|-----------|------------|------|
| Solid X Li<br>Listed as Toxic    | iquid X<br>Yes X     | Gas<br>No | ·          |      |

Status of Environmental Compliance (Past and Present):

At present, all discharges conform to existing ORNL permit conditions. The ORNL NPDES permit (1975-1980) did not contain discharge limitations for any of the listed chemicals. The present NPDES permit (1986-1990) lists only chloroform, PCBs, and trichloroethylene and these are monitoring only requirements. The primary reason for this is that existing data does not indicate that the chemicals are being released via surface water discharges. If after the mandatory year's monitoring, certain of these appear in sufficient quantity then limitations may be imposed. The organic compounds, benzene, chloroform, methylene chloride, tetrachloroethylene, and trichloroethylene appear with limits for the Nonradiological Wastewater Treatment Plant (X12), but this facility is not scheduled to be completed until 1990 and thus the limitations are not effective until that time.

Known or Suspected Public Health Impacts:

- No known or suspected public health impacts.

Known or Suspected Environmental Impacts:

The above named chemical was identified as a constituent present in spill materials that resulted in a fish-kill in White Oak Creek in 1983. See Appendix I for more details.

Level and Type of Worker Protection:

Specific use, conditions, quantities, and other factors determine the level and type of worker protection. Standard measures for protection against this material include on or more of the following: (1) adequate ventilation, (2) protective clothing, (3) chemical goggles, (4) face shield, and (5) proper respiratory protection.

Source of Chemical Usage Information:

Information taken from the Annual Toxic Chemical Usage By ORNL Department - Report 3063, ORNL Hazardous Materials Usage Reconciliation - Report 7010, and purchase requisitions with ORNL account numbers for 1980-1985.

Source of Chemical Distribution Information/Accuracy:

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories         |
|---|--|
| Other   |  |
| Off-site<br>Burial                              |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluen              |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 256,918<br>354,208<br>384,049<br>441,278<br>347,872<br>362,997 |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                           |
| Chemical<br>Name                                | Tetrachloro-<br>ethylene                                       |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

ESTIMATED CHEMICAL USAGE

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 15 t<br>8 L  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 185<br>216<br>216<br>428<br>450<br>515                 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Tetrahyodro-<br>furan                                  |

# Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 223 L<br>33 L<br>81 L                                  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 761<br>1,014<br>1,347<br>1,359<br>2,142<br>2,657       |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical                                | Toluene  |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

# Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other .                                 |  |
| Off-site<br>Burial                      | <ul><li>1 kg</li><li>5 kg</li><li>4 kg</li></ul>       |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 2,455<br>2,530<br>950                                  |
| Year                                    | 1985<br>1984<br>1983<br>1982                           |
| Chemical<br>Name                        | Tributyl<br>phosphate                                  |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

# Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| site<br>ial Other                       |  |
| On-site Off-site<br>Burial Burial Other | 14 L<br>998 L<br>26 L                                  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 | •  |
| Quantity<br>Used<br>(kg)                | 4,315<br>4,315<br>2,309<br>5,636<br>17,547             |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981<br>1980           |
| Chemical<br>Name                        | Trichloro-<br>ethylene                                 |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 48<br>2<br>5<br>2                                      |
| Year                                    | 1985<br>1984<br>1983<br>1982                           |
| Chemical<br>Name                        | Trichloro-<br>methylsilane                             |

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              | < 1 kg<br>4 kg   |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 16<br>L  |
| Year  | 1985<br>1984<br>1982<br>1981                           |
| Chemical<br>Name                                | Triethano-<br>lamine                                   |

ESTIMATED CHEMICAL USAGE

| On-site Off-site<br>Burial Burial Other Unknown | Kecords do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              | 1 Kg   |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 946<br>38  |
| Year  | 1983   |
| Chemical<br>Name                                | Urethane   |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories   |
|---|--|
| Other                                   |  |
| On-site Off-site<br>Burial Burial Other | 208 L<br>624 L<br>1,052 L                                |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| ie Liquid<br>: Effluent                 |  |
| Airborne<br>Release E                   |  |
| Product                                 | ·  |
| Quantity<br>Used<br>(kg)                | 24,129<br>22,718<br>25,617<br>29,474<br>38,686<br>40,594 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                     |
| Chemical<br>Name                        | Varsol   |

| Chemical<br>Name | Year | Quantity<br>Used<br>(kg) | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Unknown                                  |
|------------------|------|--------------------------|---------|---------------------|-------------------------------------|--------------------|---|--|
| Vydate           | 1982 | 19                       |         | -                   |                                     |                    |   | Records do<br>not exist for<br>the other |
|                  |      |                          |         |                     |                                     |                    |   | categories                               |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

#### ATTACHMENT III ORNL Surplus Chemical Sales and Transfers

#### ORNL Waste Oil Sold for Recycle

| <u>Date</u>  | Quantity (liters)  | Off-site Buyer  |
|--|--|---|
| FY 1980<br>FY 1981<br>FY 1982<br>FY 1983<br>FY 1984<br>FY 1985 | 37,850<br>39,364<br>25,738<br>8,176<br>71,468<br>7,949<br>18,168 | Petroleum Recycling Corp. Alpha Recovery Systems, Inc. Petroleum Recycling Corp. Universal Oil & Supply Co. Alpha Recovery Systems, Inc. Universal Oil & Supply Inc. Alpha Recovery Systems, Inc. |
| Total  | 208,713  |   |

All wastes oils are analyzed for PCBs, organics, and radioactivity prior to being placed in the Waste Oil Storage Tank.

#### Waste Acceptance Criteria:

| PCB             | <2 ppm                     |
|-----------------|----------------------------|
| Organics        | <50 ppm_                   |
| Gross beta      | $1.1 \times 10^{-2}$ Bq/mL |
| Gross gamma     | 1.1x10 <sup>-2</sup> Bq/mL |
| 3 <sub>H</sub>  | $1.1x10^{3}$ Bq/mL         |
| 14 <sub>C</sub> | 3x10 <sup>2</sup> Bq/mL    |

#### Surplus Chemicals Transferred to Other Government Agencies

| <u>Date</u> | <u>Chemical</u>  | Quantity   | Receiving Agency                          |
|-------------|--|--|---|
| 6/79        | Iodic acid Nickelous sulfate Potassium chromate Potassium permanganate Hardening resin Argon   | 1 kg<br>4 kg<br>9 kg<br>23 kg<br>57 g<br>6 L                         | Department of General<br>Services         |
| 6/80        | 1-Amino-8-naphthol-4- sulfonic acid Benzoic anhydride acid 1-Naphthylamine-4- sulfonic acid 1-Naphthylamine-8- sulfonic acid Benzyl acetate 2,4-Dichloro-1-Naphthol 2,3-Dichloronaphthoquinone 2,7-Dihydroxynaphthalene A-Fluoronaphthalene A-Naphthonitrile B-Naphthylamine hydrochloride A-Naphthylhydrazine hydrochloride 6-Nitroquinoline  | 1000 g 100 g 25 g 1 kg 2 kg 100 g 25 g 50 g 50 g 10 g 50 g 75 g 40 g | Comparative Animal<br>Research Laboratory |
| 7/80        | o-Cresotinic acid 2,4-Dinitro-1-naphthol-7- sulfonic acid P-Fluorobenzoic acid 8-Hydroxy-5-quinoline sulfonic acid 7-Iodo-8-hydroxyquinoline- 5-sulfonic acid A-Naphthaleneacetic acid B-Naphthoxyacetic acid B-Oxynaphtholic acid Quinaldinic acid Violuric acid 1-Amino-2-naphthol hydrochloride 2-Aminonaphthoquinone 2-Chloroquinoline 6-Chloroquinoline 5,7-Dibromo-8- hydroxyquinoline | 100 g 25 g 100 g 100 g 100 g 25   | Comparative Animal<br>Research Laboratory |

#### Surplus Chemicals Transferred to Other Government Agencies

| <u>Date</u>   | <u>Chemical</u>   | <u>Quanti</u>                                      | ty                 | Receiving Agency   |
|---------------|---|--|--------------------|--|
| 7/80<br>cont. | 2,4-Dibromo-l-naphthol 9,10-Dichloroanthracene 3,5-Dimethylpyrozole 2,7-Dinitrophenanthra- quinone 2,4-Dioxothiazolidine N-Methyl-2-quinoline 5-Nitro-l-naphthylamine 2-Nitrophenanthraquinone 8-Nitroquinoline | 25 g<br>25 g<br>25 g<br>1 g<br>25 g<br>5 g<br>10 g |                    |  |
| 8/80          | <pre>1 lot (507 items) of   miscellaneous chemicals   classified as DOT Flammab   liquid, n.o.s. and   poisonous solid, n.o.s.</pre>  | le   |                    | University of Tennessee,<br>Knoxville via the<br>Department of General<br>Services |
| 7/83          | Potassium nitrite -   | 16 1   | kg                 | Georgia Coastal Plain<br>Experimental Station,<br>Tifton, GA                       |
| 12/85         | Ethyl Alcohol   | 11 (   | drums              | University of Tennessee,<br>Knoxville via the<br>Department of General<br>Services |
| 3/86          | Hydraulic fluid,<br>non-petroleum base<br>Hydraulic fluid<br>Gulf, oil<br>Marcol, oil   | 371<br>98<br>2                                     | container<br>drums | Roane State Community<br>College via the Department<br>of General Services<br>s    |
|               | Shell Hydraulic oil<br>Isopropanol  |  | drums<br>drums     |  |

#### Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u> | Chemical  | Quantity  |
|-------------|---|---|
| 2/84        | Sulfuric acid   | 16 kg   |
| 11/84       | o-Aminophenol-p-sulfonic acid Thioglycolic acid p-Anisidine acetyl o-Toluidine acetyl Dibromopropyl alcohol Cyanoacetanilide Resorcinol monomethyl ether Furil dioxime Heptoyl chloride 2-Hydroxy-1,4-Dimethyl benzene Methyl bromobenzoate Methyl-p-aminophenol sulfate Myristoyl chloride P-Naphthol benzine o-Nitrophenetole P-Nitrophenetole P-Nitrophenylhydrazine | 5 kg 200 g 400 g 2 kg 1 kg 100 g 75 g 300 g 250 g 525 g 100 g 100 g 100 g                 |
|             | P-Nitrophenylhydrazine hydrochloride P-Nitrosophenol sodium salt Palmitoyl chloride G-Phenylpropyl bromide Phloroglucinol Propionamide Tetra-N-Butylammonium iodide Thio-B-Naphthol Toluhydroguinone P-Tolyl isocyanate Trimethylamine Trimethylene bromide Trimethylene chlorohydrin Xanthydrol o-xylose   | 100 g 125 g 25 g 100 g 150 g 200 g 400 g 50 g 250 g 250 g 250 g 200 g 20 g 275 g          |
| 5/85        | Lead carbonate Potassium bromide Sodium acetate Sodium peroxide Tantalum pentoxide Adipic acid o-Aminobenzenesulfonic acid D-10-Camphorsulfonic acid Chloroacetic acid 3,5-Dinitrobenzoic acid Iodoacetic acid B-Iodopropionic acid   | 10 kg<br>25 g<br>30 kg<br>5 kg<br>1 kg<br>125 g<br>100 g<br>2 kg<br>100 g<br>25 g<br>25 g |

#### · Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u> | Chemical                           | <u>Quantity</u> |
|-------------|------------------------------------|-----------------|
| 5/85        | Naththalene-B-sulfonic acid        | 200 g           |
| cont.       | A-Naphthoic acid                   | 25 g            |
|             | o-Toluic acid                      | 800 g           |
|             | p-Toluic acid                      | 3 kg            |
|             | Undecylenic acid                   | 500 g           |
|             | Acenaphthene                       | 200 g           |
|             | Allyl formate                      | 100 g           |
|             | m-Aminophenol                      | 200 g           |
|             | o-Aminophenol                      | 100 g           |
|             | p-Aminophenol                      | 500 g           |
|             | p-Aminophenol hydrochloride        | 200 g<br>50 g   |
|             | Tert-Amylbenzene                   | 400 g           |
|             | n-Amyl bromide                     | 100 g           |
|             | o-Anisidine                        | 100 g           |
|             | p-Anisidine<br>Anthraquinone       | 250 g           |
|             | Azoxybenzene                       | 100 g           |
|             | Benzil                             | 200 g           |
|             | Benzyl chloride                    | 2 kg            |
|             | A-Bromo-acetophenone               | 200 g           |
|             | Bromobenzene                       | 10 g            |
|             | Bromomesitylene                    | 200 g           |
|             | o-Bromophenol                      | 50 g            |
| •           | 2-Bromopyridine                    | 100 g           |
|             | o-Bromotoluene                     | 50 mL           |
|             | Tert-butylbenzene                  | 25 g            |
|             | N-Butyl bromide                    | 250 g           |
|             | Isobutyl chloride                  | 200 g           |
|             | Isocaproyl chloride                | 50 g            |
|             | Cetyl bromide                      | 200 g<br>100 g  |
|             | Cetyl iodide                       | 500 g           |
|             | o-Chloroaniline<br>p-Chloroaniline | 100 g           |
|             | A-Chloromethylnaphthalene          | 500 g           |
|             | Crotonaldehyde                     | 500 g           |
|             | Cyanoacetamide                     | 200 g           |
|             | 1,2,5,6-Dibenzanthracene           | 1 g             |
|             | 9,10-Dibromoanthracene             | 50 g            |
|             | N-Dibutylamine                     | 5 kg            |
|             | Di-n-butyl Ketone                  | 25 g            |
|             | Dicyandiamide                      | 1 kg            |
|             | Diethyl Ketone                     | 3 kg            |
|             | Dimethylbenzylamine                | 200 g           |
|             | AS-Diphenylhydrazine hydrochloride | 25 g            |
|             | Di-p-tolyl Ketone                  | 25 g            |
|             | Petroleum ether                    | 8 L<br>100 g    |
|             | Ethyl anisate                      | 100 g           |
|             | Ethyl anthranilate                 | 100 g           |
|             | Ethyl B-Bromopropionate            | 100 9           |

#### Surplus Chemicals Transferred to ORGDP Property Sales

| <u>Date</u>   | Chemical                                      | Quantity       |
|---------------|---|----------------|
| 5/85<br>cont. | Ethylene chlorobromide<br>Ethylene chloride   | 500 g<br>8 kg  |
|               | Ethyl fumarate                                | 1 kg           |
|               | Ethyl maleate                                 | 2 kg           |
|               | Ethyl orthoformate                            | 250 g          |
|               | Lauryl bromide                                | 700 g          |
|               | Methyl anthranilate                           | 3 kg           |
|               | Methyl bromoacetate                           | 400 g          |
|               | Methyl N-butylcarbinol                        | 25 g           |
|               | Methyl N-butyrate                             | 200 g          |
|               | o-Methylhydroxylamine hydrochloride           | 100 g          |
|               | Methyl malonate                               | 500 g          |
|               | Methyl B-naphthyl Ketone                      | 200 g<br>600 g |
|               | Myristyl bromide                              | 100 g          |
|               | m-Nitroaniline                                | 200 g          |
|               | o-Nitroaniline                                | 200 g<br>100 g |
|               | 4-Nitro-1,2-Diaminobenzene                    | 25 g           |
|               | p-Nitrofluorobenzene                          | 100 g          |
|               | Nitron  | 100 g          |
|               | <pre>1-Nitroso-2-napthol o-Nitrotoluene</pre> | 3 kg           |
|               | * ***   | 250 g          |
|               | p-Nitrotoluene<br>Phenanthraquinone           | 25 g           |
|               | Phenetole                                     | 500 g          |
|               | Phenylbenzylcarbinol                          | 25 g           |
|               | m-Phenylenediamine                            | 250 g          |
|               | DL-A-Phenylethylamine                         | 100 g          |
|               | DL-Phenylmethylcarbinol                       | 5 kg           |
|               | Picoline                                      | 500 g          |
|               | Propiophenone                                 | 1000 g         |
|               | Iso-propylamine                               | 8 kg           |
|               | Iso-propyl iodide                             | 100 g          |
|               | Quinone                                       | 2 kg           |
|               | Trans-Stilbene                                | 300 g          |
|               | Strychnine sulfate                            | 200 g          |
|               | Succinonitrile                                | 200 g          |
|               | Sucrose                                       | 150 g          |
|               | m-Tolunitrile                                 | 25 g           |
|               | o-Tolunitrile                                 | 25 g           |
|               | p-Tolunitrile                                 | 100 g          |
|               | N-Valeronitrile                               | 100 g          |
|               | Bromocresol green                             | 5 L            |
|               | Buffer solution                               | 64 L           |

| <u>Date</u> | Chemical  | <u>Quantity</u>  |
|-------------|---|--|
| 6/80        | 4-Iodo-n-xylene   | 250 g  |
| 6/80 8/81   | Silicic acid Sulfuric acid Ammonium thiocyanate Calcium chloride Sodium methylate Yttrium, metal 2-Amino-5-chlorobenzoic acid 1-Amino-2-naphthol-4-sulfonic acid o-Aminophenol-p-sulfonic acid m-Chlorobenzoic acid M-Chlorobenzoic acid Chlorosulfonic acid 3-Hydroxy-2-naphthoic acid 1-Malic acid DL-Mandelic acid Methoxyacetic acid o-Methoxybenzoic acid 3-Nitrosalicylic acid 5-Nitrosalicylic acid Salicylacetic acid Salicylacetic acid Salicylacetic acid Salicylacetic acid Acetamide Allylthiourea N-Amyl chloride Tert-Amyl chloride Benzyl bromide Benzyl bromide Benzyl bromide O-Bromotoluene sec-Butyl bromide N-Butyl-p-toluenesulfonate Chloroacetdiethylamide Chloroacetonitrile Chloroacetyl chloride A-Chloronaphthalene 2-Chloropyridine Cinchonine, alkaloid o-Cresyl-p-toluenesulfonate Dicyclohexylamine Ethyl bromo-n-butyrate | 2332742724100120100013125001000130013001300130013001300130013001 |
|             | Ethyl bromo-n-caproate Ethyl chloroacetate Ethyl naphthylamine m-Fluorochlorobenzene  | 75 g<br>250 g<br>100 g<br>25 g<br>25 g                           |
|             | m-Fluorotoluene<br>o-Fluorotoluene  | 100 g  |

| <u>Date</u>  | <u>Chemical</u>   | <u>Quantity</u>  |
|--------------|---|--|
| 8/81<br>cont | 2-Iodo-1,3-dimethylbenzene Methyl adipate Methyl bromopropionate Methyl orthoacetate 6-Methylquinoline 8-Methylquinoline Methyl toluate Naphthyl acetate p-Nitroacetanilide A-Nitronaphthalene m-Nitrophenol Phenylethyl bromide Phenylhydrazine DL-Phenylmethylcarbinol Phenyl-p-tolyl ketone Pinacol Propiophenone N-propyl carbonate Propylene glycol Thionaphthol Ethyl bromopropionate Glycine anhydride   | 10 g<br>100 g<br>300 g<br>600 g<br>25 g<br>50 g<br>75 g<br>400 g<br>25 g<br>300 g<br>500 g<br>500 g<br>500 g<br>100 g<br>100 g<br>75 g             |
| 7/82         | Hydrofluosilicic acid Cadmium metal Calcium chloride Lead sulfate Magnesium sulfate Potassium thiocyanate A-Bromo-n-valeric acid o-Chlorobenzoic acid Trans-Cinnamic acid 2,4-Dinitrobenzene sulfonic acid Mucic acid Naphthalic anhydride acid m-Toluic acid. Trichloroacetic acid Diacetone alcohol Myristyl alcohol Phenylethyl alcohol Phenylpropyl alcohol 2-Aminopyrimidine Iso-amyl bromide N-Amyl iodide Benzalacetone Benzidine dihydrochloride Bromoacetyl bromide Bromocyclohexane N-Butyl bromide | 3 kg 113 g 11 kg 16 kg 16 kg 100 g 100 g 100 g 5 kg 100 g 5 kg 250 g 100 g 300 g 100 g 300 g 15 kg |

| <u>Date</u>     | <u>Chemical</u>  | <u>Quantity</u>  |
|-----------------|--|--|
| Date 7/82 cont. | N-Butyl sebacate Cellulose acetate Cellulose triacetate m-Chlorophenol o-Chlorophenol Di-n-butylaniline p-Diethylaminobenzaldehyde Diethyl-A-naphthylamine Ethylamine hydrochloride Ethyl bromoacetate Ethylene bromide Ethylene chlorohydrin Ethyl phosphate Fluoranthene Fluorenone N-Hexylphenylcarbinol  | 500 g<br>1 kg<br>500 g<br>25 g<br>1 kg<br>300 g<br>25 g<br>25 g<br>200 g<br>900 g<br>23 L<br>2 kg<br>200 g<br>100 g<br>40 g  |
|                 | N-Hexylphenylcarbinol o-Idobenzoyl chloride Methyl N-amyl Ketone Methylethylamine hydrochloride 4-Methyl-2-pentanol Methyl phenylacetate Methyl-iso-thiocyanate 4-Methylumbelliferone 1,2-Naphthoquinone 1-Naphthyl isocyanate Pentaerythritol Phenanthrene m-Phenetidine o-Phenetidine p-Phenetidine Phenyl acetate Phenyl-a-naphthylamine Phthalyl chloride p-Isopropylbenzaldehyde Isopropyl bromide Propylene chloride N-Propyl sulfone p,p-Tetramethyldiaminodiphenylmethane o-Tolidine dihydrochloride 1,2,3-Tribromopropane 1,1,1-Trichloroethane | 100 g<br>100 g<br>60 g<br>50 g<br>250 g<br>25 g<br>25 g<br>25 g<br>100 g |
| 8/82            | Xanthydrol  1,1,1-Trichloroethane 2,2,4-Dihydroxybenzaldehyde  | 80 g<br>416 L<br>700 g   |
| 4/83            | Ammonium acetate<br>Ammonium sulfide<br>Antimony, metal  | 16 kg<br>5 L<br>< 1 kg   |

| <u>Date</u> | <u>Chemical</u>  | <u>Quantity</u>  |
|-------------|--|--|
| 4/83 cont.  | Calcium fluoride Ferrous sulphamate Lithium sulfate Mercurous nitrate Acetyl p-aminobenzoic acid B-Amino-l-naphthol-3,6-disulfonic acid DL-A-Aminophenylacetic acid 4-Amino-m-toluenesulfonic acid p-Arsanilic acid DL-10-Camphorsulfonic acid Cyanuric acid A,B-Dibromosuccinic acid Diglycolic acid Furoic acid Indole-3-acetic acid Acetyl thiourea B-Diethylaminoethyl alcohol 1-Aminoanthraquinone Benzoylacetone Benzyl thiocyanate Bromohydroquinone N-Butyl sulfone 2,4-Dibromophenol Di-isobutylene Dichloroacetyl chloride 1,3-Diphenylguanidine Ethyl,N-Butylmalonate Ethyl isothiocyanate Hexachloroethane Methyl thiocyanate Phenyl isothiocyanate Phenyl isothiocyanate Propylene diamine Succinimide p,p-Tetramethyldiaminobenzophenone Xanthone Ion Exchange Resin | 5 L g l kg                 |
| 4/84        | Molybdic acid Cesium oxide Potassium bifluoride Potassium, metal Potassium periodate Potassium thiocyanate Titanium oxide 1-Nitroanthraquinone-5-sulfonic acid Trichloroacetic acid Acetamide p-Aminophenol acetyl B-Dimethylaminoethyl alcohol  | 1 kg 5 kg 5 kg 1 kg 10 kg 10 kg 5 g 800 g 1 kg 25 g 1 kg |

| <u>Date</u>   | Chemical   | <u>Quantity</u>  |
|---------------|--|--|
| 4/84<br>cont. | p-Aminoacetophenone 2-Aminoanthraquinone 3-Aminoquinoline 2-Aminothiazole Benzenesulfonylamide o-Bromophenetole p-Bromophenetole | 50 g<br>100 g<br>50 g<br>1000 g<br>100 g<br>125 g<br>100 g |
|               | m-Chlorophenyl isocyanate  | 100 g  |
|               | 2,4-Diaminophenol dihydrochloride Dibenzalacetone  | 500 g<br>25 g  |
|               | 4,4-Dibromobiphenyl  | 200 g  |
|               | 2-Diethylamino-1,4-dimethylbenzene   | 50 g<br>50 g   |
|               | 4,4-Dihydroxybenzophenone 2,5-Dimethoxyaniline   | 75 g   |
|               | Dimethyldihydroresorcinol  | 25 g   |
|               | Phenyl cellosolve  | 22 kg  |
|               | Ethylene glycol, diacetate   | 8 kg   |
|               | Hydantoin  | 75 g<br>50 g   |
|               | a-Hydrindone   | 200 g  |
|               | p-Hydroxybenzaldehyde<br>g-Hydroxybutronitrile   | 10 g   |
|               | p-Hydroxyphenylglycine   | 100 g  |
|               | o-Methoxybenzaldehyde  | 300 g  |
|               | p-Phenylacetophenone   | 25 g   |
|               | Isopropyl acetate  | 100 g  |
|               | Isopropyl acetate, purified  | 8 L.   |
|               | Tetramethylammonium chloride   | 1000 g   |
|               | p-Toluamide  | 75 g<br>200 g  |
|               | 1,2,3-Trichloropropane   | 200 g<br>23 L  |
|               | Buffer solution  | 23 L<br>8 kg   |
|               | Ion Exchange resin   | 0 78   |

#### APPENDIX I

On August 6, 1983, an unknown quantity of organic mixtures and pesticides spilled into White Oak Creek resulting in a fish-kill. Approximately 7,570 L of material (mostly water) was removed from the creek and shipped to an off-site facility for disposal. The chemical analyses of this material is shown below.

#### ANALYSES RESULTS OF WHITF OAK CREEK FISH KILL INCIDENT

| Name of Chemical   | Concentration   |
|--------------------|-----------------|
| Aldrin             | Trace (low ppb) |
| Heptachlor         | Trace (low ppb) |
| Acetaldehyde       | 1 ppm           |
| Freon 113          | 1 ppm ·         |
| Xylene             | < 0.5 ppm       |
| Toluene            | < 0.5 ppm       |
| Freon 11           | < 0.5 ppm       |
| Benzene            | < 0.5 ppm .     |
| Ethyl alcohol      | < 0.5 ppm       |
| Methylene chloride | < 0.5 ppm       |
| Acetone            | < 0.5 ppm       |
| Butyl cellusolve   | < 0.5 ppm       |
| Dimethyl benzene   | < 015 ppm       |

History of PCB-Related Incidents/Spills at ORNL

APPENDIX II

| <u>Date</u>        | <u>Location</u> | <u>Material</u>                           | Amount                |
|--------------------|-----------------|---|-----------------------|
| May 6, 1981        | 3026            | PCB Oil<br>(Transformer)<br>< 5 ppm       | small<br>amount       |
| July 27, 1981      | 3012            | PCB 0il                                   | 378.5 L<br>(100 gal)  |
| August 3, 1981     | 6000            | PCB Oil                                   | 3.78 L<br>(1 gal)     |
| August 12, 1983    | 4500S, R-211    | PCB Oil<br>(light ballast)<br>480,000 ppm | 0.01 L<br>(0.003 gal) |
| April 4, 1984      | 3026-D          | PCB Oil<br>(Transformer)<br>500 ppm       | 0.47 L<br>(1 pint)    |
| April 16, 1986     | 4500S, D-61     | PCB Oil<br>(light ballast)<br>854 ppm     | 0.01 L<br>(0.003 gal) |
| September 24, 1985 | 4500S, G-260    | PCB Oil<br>(light ballast)<br>930,000 ppm | 0.01 L<br>(0.003 gal) |

#### ATTACHMENT II

Estimated Chemical Usage

| Unknown   | Records do<br>not exist<br>for the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site Of<br>Burial Bu                         | 35 L<br>61 L<br>36 L                                   |
| On-site C<br>Storage B                          |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 10,447<br>12,858<br>13,278<br>15,473<br>16,858         |
| Year  | 1985<br>1984<br>1983<br>1982<br>1980                   |
| Chemical<br>Name                                | Acetone  |

<sup>\*\*&</sup>quot;Off-site burial" does not necessarily imply that the wastes were managed by land based disposal. Treatment methods (incineration) has been utilized in some cases. This point is applicable throughout this report.

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managea as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 12 L<br>24 L<br>13 L                                   |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 25,204<br>31,162<br>26,930<br>18,501<br>14,020         |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981<br>1980           |
| Chemical<br>Name                                | Acetonitrile   |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(L) | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site Off<br>Burial Bu | On-site Off-site<br>Burial Burial Other | Unknown  |
|------------------|--------------------------------------|-------------------------|---------|---------------------|-------------------------------------|--------------------|--------------------------|---|--|
| Acrylamide       | 1985<br>1984<br>1983<br>1982<br>1981 | 2<br>5<br>4<br>1,368    |         |                     |                                     |                    | .5 kg<br>3 kg<br>9 kg    | к <u>9</u><br>9                         | Records do<br>not exist for<br>the otner<br>categories |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      | 324 kg<br>117 kg<br>6 kg                               |
| On-site Off-site<br>Burial Burial Other | in   |
| On-site<br>Storage                      | 2, 479 L<br>Presently in<br>storage                    |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 3,818<br>3,636   |
| Year                                    | 1985<br>1984<br>1983                                   |
| Chemical<br>Name                        | Aluminum nitrate                                       |

| Unknown                                 | Records do<br>exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 3,330<br>3,391<br>5,993<br>4,232<br>3,634<br>6,709 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981               |
| Chemical<br>Name                        | Ammonia,<br>anhydrous                              |

ESTIMATED CHEMICAL USAGE

| Unknown   | Records do<br>not exist<br>for tne other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 6 kg<br>< 1 kg<br>< 1 kg                               |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 15<br>69<br>70<br>198<br>305<br>657                    |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Ammonium<br>bifluoride                                 |

| Chemical<br>Name      | Year                                 | Quantity<br>Used<br>(L)                            | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site Off<br>Burial Bu | On-site Off-site<br>Burial Burial Other | Unknown  |
|-----------------------|--------------------------------------|--|---------|-------------------------------------|--------------------|--------------------|--------------------------|---|--|
| Ammonium<br>hydroxide | 1985<br>1984<br>1983<br>1982<br>1981 | 1,135<br>1,524<br>2,053<br>1,666<br>1,540<br>2,549 |         |                                     |                    |                    | 28<br>277<br>309         | ку<br>8<br>8                            | Records do<br>not exist for<br>the other<br>categories |

ESTIMATED CHEMICAL USAGE

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 4 kg<br>4 kg<br>2 kg                                   |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 281<br>255<br>200<br>826<br>270<br>2,613               |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Ammonium<br>nitrate                                    |

| Chemical<br>Name                 | Year                                 | Quantity<br>Used<br>(L)   | Product | Airborne<br>Release | Liquid<br>Effluent | On-site<br>Storage              | On-site Off-site<br>Burial Burial Other   | Unknown |
|----------------------------------|--------------------------------------|---|---------|---------------------|--------------------|---------------------------------|---|---------|
| Asbestos-Containing<br>Materials | ug                                   |   |         |                     |                    |                                 |   |         |
| - Felt Roofing                   | 1985<br>1984<br>1983<br>1982<br>1981 | 972m2<br>640m2<br>126m2<br>1,655m2<br>2,173m2<br>3,060m2  |         |                     |                    | 1985 2<br>1<br>1984 2<br>1983 2 | 241 m <sup>2</sup> On-site burial 1,426 m On-site burial 28 m <sup>2</sup> On-site burial 2,982 m On-site burial 274 m <sup>2</sup> On-site burial 1,954 m On-site burial |         |
| - Sheet Roofing                  | 1985<br>1984<br>1983<br>1982<br>1981 | 1,795m <sup>2</sup><br>1,541m <sup>2</sup><br>1,463m <sup>2</sup><br>702m <sup>2</sup><br>488m <sup>2</sup> |         |                     |                    |                                 |   |         |
| - Tape                           | 1984<br>1983<br>1982<br>1981<br>1980 | 56 rolls<br>48 rolls<br>15 rolls<br>38 rolls<br>89 rolls  | ຑຑຑຑຑ   |                     |                    |                                 |   | ·       |

<sup>\*</sup>There are unknown quantities of asbestos-containing insulation throughout ORNL. A major effort has been made during the last 3-4 years to identify these areas, remove the asbestos-containing insulation, dispose of properly, and replace with nonasbestos insulation.

| tity<br>ed Airborne Liquid On-site On-site Off-site<br>) product Release Effluent Storage Burial Burial Other Unknown<br>) | (cont.)                               | 3m<br>5m<br>9m<br>5m<br>6m<br>1m                | 3 ea<br>9 ea<br>6 ea<br>8 ea<br>7 ea<br>0 ea                         | 4 <sup>m</sup> <sup>2</sup><br>6 kg | 0m <sup>2</sup><br>0 kg     | 9m <sup>2</sup><br>4 kg    | $9m^2$                     | $4m^2$ 5 kg .                 | c     |
|--|---------------------------------------|---|--|-------------------------------------|-----------------------------|----------------------------|----------------------------|-------------------------------|-------|
| Quantity<br>Used<br>(L) Produc   | ls (cont.)                            | 53m<br>295m<br>189m<br>165m<br>456m             | 3,193 ea<br>3,669 ea<br>3,396 ea<br>2,408 ea<br>2,337 ea<br>2,450 ea | 264m <sup>2</sup><br>166 kg         | 270m <sup>2</sup><br>220 kg | 319m <sup>2</sup><br>94 kg | 319m <sup>2</sup><br>94 kg | 384m <sup>2</sup><br>155 kg · | 2.010 |
| Qu<br>Year   | ining Materia                         | re 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | 1985 3<br>1984 3<br>1983 3<br>1982 2<br>1981 2                       | 1985                                | 1984                        | 1983                       | 1982                       | 1981                          | 0     |
| Chemical<br>Name   | Asbestos-Containing Materials (cont.) | - Insulated Wire                                | - Gaskets  | - Packing                           |                             |                            |                            |                               |       |

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
|   |  |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| /<br>Product                                    | ·  |
| Quantity<br>Used<br>(L)                         | 1,918  |
| Year  | 1981   |
| Chemical<br>Name                                | Barium<br>octahydrate                                  |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| 0ther   |  |
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 15   |
| Year  | 1984   |
| Chemical<br>Name                                | Baygon (N-Methyl 1984<br>carbamate)                    |

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(L)         | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other Unknown | Unknown  |
|------------------|--------------------------------------|---------------------------------|---------|---------------------|-------------------------------------|--------------------|-------------------|---|--|
| Benzene          | 1985<br>1984<br>1983<br>1982<br>1981 | 238<br>110<br>246<br>530<br>708 |         | ·                   |                                     |                    |                   | 37 L<br>43 L<br>38 L                            | Records do<br>not exist for<br>the other<br>categories |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

ESTIMATED CHEMICAL USAGE

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      |  |
| On-site Off-site<br>Burial Burial Other | <ul><li>1 Kg</li><li>1 Kg</li><li>1 Kg</li></ul>       |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne<br>Release E1                  |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 16<br>19   |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Cadmium<br>nitrate                                     |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 8 kg<br>5 kg   |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 1,216<br>936<br>952<br>1,095<br>1,772<br>1,325         |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Calcium<br>hypochlorite                                |

ESTIMATED CHEMICAL USAGE

| Chemical<br>Name | Year | Quantity<br>Used<br>(L) | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other Unknown | te<br>1 Other | Unknown       |
|------------------|------|-------------------------|---------|-------------------------------------|--------------------|--------------------|---|---------------|---------------|
| Carbon           | 1985 | 13                      |         |                                     |                    |                    |   |               | Records do    |
| monoxide         | 1984 | 27                      |         |                                     |                    |                    |   |               | not exist for |
|                  | 1983 | 54                      |         |                                     |                    |                    |   |               | rile oriler   |
|                  | 1982 | 29                      |         |                                     |                    |                    |   |               | caredories    |
|                  | 1981 | 63                      |         |                                     |                    |                    |   |               |               |
|                  | 1980 | 20                      |         |                                     |                    |                    |   |               |               |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 29 L<br>64 L   |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             | -  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 117<br>273<br>204<br>208<br>341<br>356                 |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Carbon<br>tetrachloride                                |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Unknown                                 |  |   |
|---|--|---|
| 0ther                                   |  |   |
| On-site Off-site<br>Burial Burial Other |  |   |
| On-site<br>Burial                       |  |   |
| On-site<br>Storage                      |  |   |
| Liquid<br>Effluent                      |  | e e g   |
| Airborne<br>Release                     |  | Approximately 1590 kg<br>is used to treat the<br>the Sewage Treatment<br>Plant effluent |
| Product                                 |  | Approxi<br>is used<br>the Sew<br>Plant e  |
| Quantity<br>Used<br>(L)                 | 3,772<br>3,363<br>3,568<br>2,681<br>2,554<br>3,295 | 1,590/yr.   |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981<br>1981       | 1950-1979   |
| Chemical<br>Name                        | Chlorine,<br>gas                                   |   |

### Estimated Distribution in %

| Chemical<br>Name | Year   | Quantity<br>Used<br>(L)                | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other Unknown | . Unknown  |
|------------------|--|--|---------|---------------------|-------------------------------------|--------------------|-------------------|---|--|
| Chloroform       | 1985<br>1984<br>1983<br>1982<br>1981<br>1980 | 159<br>121<br>114<br>246<br>261<br>167 |         |                     | •                                   |                    |                   | 19 L<br>37 L<br>68 L                            | Records do<br>not exist for<br>the other<br>categories |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| nknown  | Records do<br>not exist for<br>the other |
|---|--|
| ther U  | X 5 7 0                                  |
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site (<br>Burial                             |  |
| On-site O<br>t Storage B                        |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 76<br>61                                 |
| Year  | 1982<br>1981                             |
| Chemical<br>Name                                | Chlorpyrifos<br>(Bolt)                   |

ESTIMATED CHEMICAL USAGE

| 606 L Records do 77 L not exist for 188 L the other categories | i i   |
|--|-------|
|  | 886   |
|  | 787   |
| categories   | 1.911 |
|  | 806   |
|  | 2,502 |
|  | 1,590 |

ESTIMATED CHEMICAL USAGE

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown |  |
| e Off-<br>Bur                                   |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 18,925   |
| Year  | 1982   |
| Chemical<br>Name                                | Coal-oil mixture 1982                                  |

ESTIMATED CHEMICAL USAGE

| Chemical<br>Name | Year                 | Quantity<br>Used<br>(L)  | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | 0ther | Unknown  |
|------------------|----------------------|--------------------------|---------|---------------------|-------------------------------------|--------------------|---|-------|--|
| Coal Tar Pitch   | 1984<br>1983<br>1981 | 15,405<br>4,164<br>2,445 |         |                     |                                     |                    | 103 L                                   |       | Records do<br>not exist for<br>the other<br>categories |

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(L)      | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | Unknown  |
|------------------|--------------------------------------|------------------------------|---------|-------------------------------------|--------------------|--------------------|-------------------|---|--|
| Cyclohexane      | 1985<br>1984<br>1983<br>1982<br>1981 | 30<br>64<br>61<br>102<br>129 |         |                                     |                    |                    |                   | 21 L<br>20 L<br>8 L                     | Records do<br>not exist for<br>the other<br>categories |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| On-site<br>Storage                                |
|---|
| Airborne Liquid On-sit<br>Release Effluent Storag |

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| te<br>1 Other                                   |  |
| Off-si<br>Buria                                 |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             | -  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 189  |
| Year  | oxy 1983   |
| Chemical<br>Name                                | 2,4-Dichloropenoxy 1983 acetic acid                    |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 95 kg  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 3 2,182  |
| Year                                    | 1985<br>1984<br>1983                                   |
| Chemical<br>Name                        | Diethylbenzene   |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

ESTIMATED CHEMICAL USAGE

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 545  |
| Year                                    | 1984   |
| Chemical<br>Name                        | Diethylenetria-<br>mine pentacetic<br>sodium salt      |

ESTIMATED CHEMICAL USAGE

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 20 L<br>77 L<br>80 L                                   |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      | -  |
| Airborne Liquid<br>Release Effluent     | -  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 8<br>45<br>23<br>61<br>19                              |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | l,4-Dioxane  |

| Unknown   | Kecords do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site On-<br>Storage Bur                      | ·  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| /<br>Product                                    |  |
| Quantity<br>Used<br>(L)                         | 5  |
| Year  | 1982   |
| Chemical<br>Name                                | Disulfoton<br>(Disyston G)                             |

| ,457<br>,673<br>11 | 1984 1,457<br>1983 1,673<br>1982 11 |
|--------------------|-------------------------------------|

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories   |
|---|--|
| On-site Off-site<br>Burial Burial Other | ìn   |
| On-site<br>Storage                      | 15,540 L<br>Presently in<br>storage  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 | and the state of t |
| Quantity<br>Used<br>(L)                 | 5,413<br>5,413   |
| Year                                    | 1984   |
| Chemical<br>Name                        | Endcor   |

| Chemical<br>Name | Year                         | Quantity<br>Used<br>(L) | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | Other | Unknown  |
|------------------|------------------------------|-------------------------|---------|-------------------------------------|--------------------|--------------------|-------------------|---|-------|--|
| Epoxy Resins     | 1985<br>1984<br>1983<br>1983 | 379<br>36<br>19<br>19   |         |                                     |                    |                    |                   | 19 L<br>69 L<br>235 L                   |       | Records do<br>not exist for<br>the other<br>categories |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other |  |
| Off-site<br>Burial                      | 3 F<br>19 F<br>91 F                                    |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      | •  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 261<br>322<br>435<br>250<br>231<br>288                 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Ethyl acetate  |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 5 L<br>17 L<br>45 L                                    |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             | •  |
| Product   | •  |
| Quantity<br>Used<br>(L)                         | 4,648<br>4,444<br>4,943<br>8,017<br>12,438             |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Ethyl alcohol  |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              |  |
| On-site<br>Burial                               |  |
| On-site O<br>Storage B                          |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 2,354<br>3,164<br>3,679<br>1,980<br>2,048<br>1,858     |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Ethyl alcohol,<br>denatured                            |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(L)                | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site<br>Burial | Off-site<br>Burial Oth | Burial Burial Other Unknown                            |
|------------------|--------------------------------------|--|---------|-------------------------------------|--------------------|--------------------|-------------------|------------------------|--|
| Ethyl ether      | 1985<br>1984<br>1983<br>1982<br>1981 | 305<br>495<br>455<br>745<br>707<br>355 |         |                                     |                    | _                  |                   | 6 kg<br>30 kg<br>36 kg | Records do<br>not exist for<br>the other<br>categories |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (D001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| er Unknown                                      | Records do<br>not exist for<br>the other<br>categories   |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 5 L<br>1488 L<br>900 L                                   |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             | ٠  |
| Airborne<br>Release                             | -  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 16,484<br>18,683<br>27,392<br>18,895<br>18,679<br>14,296 |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                     |
| Chemical<br>Name                                | Ethylene glycol  |

| u,                                      | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Unknown                                 | Records do<br>not exist f<br>the other<br>categories   |
| Other                                   |  |
| Off-site<br>Burial                      |  |
| On-site Off-site<br>Burial Burial Other | •  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 9<br>41<br>27<br>23                                    |
| Year                                    | 1985<br>1984<br>1983<br>1981                           |
| Chemical<br>Name                        | Ethylene oxide   |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      | < 1 kg<br>48 kg<br>11 kg                               |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 | ·  |
| Quantity<br>Used<br>(L)                 | 705<br>216<br>216<br>628<br>894<br>1,227<br>1,255      |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Ferric chloride  |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| e<br>Other                              |  |
| On-site Off-site<br>Burial Burial Other | < 1 kg<br>95 kg<br>< 1 kg                              |
| On-site<br>Burial                       | in '   |
| On-site<br>Storage                      | ll36 L<br>Presently in<br>storage                      |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 1,000  |
| Year                                    | 1985<br>1984<br>1983<br>1981                           |
| Chemical<br>Name                        | Ferric sulfate   |

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(L)              | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other Unknown | Other | Unknown  |
|------------------|--------------------------------------|--------------------------------------|---------|---------------------|-------------------------------------|--------------------|-------------------|---|-------|--|
| Formaldehyde     | 1985<br>1984<br>1983<br>1982<br>1981 | 503<br>284<br>47<br>121<br>121<br>98 |         |                     |                                     |                    |                   | 3 L<br>21 L<br>9 L                              |       | Records do<br>not exist for<br>the other<br>categories |

ESTIMATED CHEMICAL USAGE

|          | Un-site Uit-site<br>Burial Burial Other Unknown | Records do | not exist for | the other | categories |      |  |
|----------|---|------------|---------------|-----------|------------|------|--|
|          | Other   |            |               |           |            |      |  |
| +:0 330  | Burial  | 10 L       | <u>ک</u><br>د | 9 L       |            |      |  |
|          | Un-site<br>Burial                               |            |               |           |            |      |  |
|          | Un-site<br>Storage                              |            | -             |           |            |      |  |
| :        | Airborne Liquid<br>Release Effluent             |            |               |           |            |      |  |
|          | Airborne<br>Release                             |            |               |           |            |      |  |
|          | Product   |            |               |           |            |      |  |
| Quantity | Used<br>(L)                                     | 4          | 34            | &         | 4 16       | ω    |  |
|          | Year  | 1985       | 1984          | 1983      | 1982       | 1981 |  |
|          | Chemical<br>Name                                | Formamide  |               |           |            |      |  |

| u wo                                    | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Unknown                                 | Recc<br>not<br>the<br>cate                             |
| On-site Off-site<br>Burial Burial Other | 39 L<br>41 L<br>52 L                                   |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     | -  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 628<br>1,272<br>1,014<br>1,313<br>1,075                |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Hexane   |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (0001). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| te On-site Off-site<br>ge Burial Burial Other Unknown | 25 L Records do<br>73 L not exist for<br>123 L the other<br>categories |
|---|--|
| On-site<br>Storage                                    |  |
| Liquid<br>Effluent                                    |  |
| Airborne<br>Release E                                 |  |
| Product   |  |
| Quantity<br>Used<br>(L)                               | 669,415<br>647,076<br>843,888<br>219,295<br>120,564                    |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                                   |
| Chemical<br>Name                                      | Hydrochloric<br>acid   |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 5 kg<br>26 kg<br>2 kg                                  |
|   | 5<br>26<br>2   |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(L)                         | 921<br>1,230<br>1,262<br>940<br>1,402                  |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Hydrofluoric<br>acid                                   |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| On-site Off-site<br>Burial Burial Other | 7 kg<br>1 kg<br>13 kg                                  |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(L)                 | 1,341<br>2,020<br>1,884<br>2,137<br>1,785<br>3,296     |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981<br>1980           |
| Chemical<br>Name                        | Hydrogen<br>peroxide                                   |

| On-site Off-site<br>Burial Burial Other Unknown | Records do | not exist for | the other | categories |
|---|------------|---------------|-----------|------------|
| On-site Off-site<br>Burial Burial               |            |               |           |            |
| On-site<br>Storage                              |            |               |           | •          |
| Airborne Liquid<br>Release Effluent             |            |               |           |            |
| Airborne<br>Release                             |            |               |           |            |
| Product   |            |               |           | •          |
| Quantity<br>Used<br>(L)                         | •          | 409           |           |            |
| Year  |            | 1985          |           |            |
| Chemical<br>Name                                |            | Isopentane    |           |            |

## Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories  |
|---|---|
| Other                                   |   |
| On-site Off-site<br>Burial Burial Other | 628 L<br>4 L  |
| On-site<br>Burial                       |   |
| On-site<br>Storage                      |   |
| Liquid<br>Effluent                      |   |
| Airborne<br>Release                     |   |
| Product                                 | ·   |
| Quantity<br>Used<br>(L)                 | 5,988<br>10,628<br>13,263<br>14,599<br>13,963<br>20,787 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                    |
| Chemical<br>Name                        | Isopropyl<br>alcohol                                    |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Chemical<br>Name  | Year | Quantity<br>Used<br>(L) | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Unknown  |
|---|------|-------------------------|---------|---------------------|-------------------------------------|--------------------|---|--|
| Kelthane<br>(4,4'-Dichloro-<br>α-trichloromethyl<br>benzhydrol) | 1985 | 61                      |         |                     |                                     |                    | 7 L                                     | Records do<br>not exist for<br>the other<br>categories |

|                    |                                      |   |         |                                     |                    | :                  |                   |   |  |
|--------------------|--------------------------------------|---|---------|-------------------------------------|--------------------|--------------------|-------------------|---|--|
| Chemical<br>Name   | Year                                 | Quantity<br>Used<br>(L)                   | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | Unknown  |
| Lacquer<br>thinner | 1985<br>1984<br>1983<br>1982<br>1981 | 6,226<br>6,942<br>7,684<br>8,123<br>9,459 |         |                                     |                    |                    |                   | 2 L<br>19 L                             | kecords do<br>not exist for<br>the other<br>categories |

|   |         |                         |         |                                     | •                  |                    |   |               |
|---|---------|-------------------------|---------|-------------------------------------|--------------------|--------------------|---|---------------|
| Chemical<br>Name                                | Year    | Quantity<br>Used<br>(L) | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | on-site<br>Storage | On-site Off-site<br>Burial Burial Other Unknown | . Unknown     |
|   |         |                         |         |                                     |                    |                    |   |               |
|   |         | (                       |         |                                     |                    |                    |   | Records do    |
| Lasso   | 1982    | ה                       |         |                                     |                    |                    |   | not exist for |
| (2-Chloro-2',6'-                                |         |                         |         |                                     |                    |                    |   | the other     |
| <pre>diethyl-N-Methoxymethyl acetanilide)</pre> | ymethyl |                         |         |                                     |                    |                    |   | categories    |
|   |         |                         |         |                                     |                    |                    |   |               |

| Unknown                                 | Records do<br>not exist<br>for the other<br>categories |
|---|--|
| Other                                   |  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne<br>Release                     |  |
| Product                                 | ·  |
| Quantity<br>Used<br>(kg)                | 3,238<br>4,934<br>6,892<br>6,095<br>3,920<br>5,471     |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Lead, sheet  |

| Unknown   |      | Records do | not exist for | the other | categories |
|---|------|------------|---------------|-----------|------------|
| On-site Off-site<br>Burial Burial Other Unknown |      |            |               |           |            |
| On-si<br>Buria                                  |      |            |               |           |            |
| On-site<br>Storage                              | ,    |            |               |           |            |
| Airborne Liquid<br>Release Fffluent             |      |            |               |           |            |
| Airborne  | 3637 |            |               |           |            |
| + C = PC × Q                                    | _    |            |               |           |            |
| Quantity<br>Used                                | (KB) | ,          |               |           |            |
| ,   | rear |            | 1983          |           |            |
| Chemical  | Name |            | Lindane       |           |            |

|  | 1985 91 · 1984 727 1982 12 |
|--|----------------------------|

| e On-site Off-site<br>e Burial Burial Other Unknown | Records do not exist for the other categories |
|---|---|
| On-site<br>Storage                                  |   |
| Liquid<br>Effluent                                  |   |
| Airborne<br>Release                                 |   |
| Product   | ·   |
| Quantity<br>Used<br>(kg)                            | 76<br>76<br>91                                |
| Year  | 1983<br>1982<br>1981                          |
| Chemical<br>Name                                    | Malathion                                     |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 225 kg<br>43 kg<br>75 kg                               |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     | •  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 97<br>139<br>322<br>250<br>304<br>576                  |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Mercury  |

## Estimated Distribution in %

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories       |
|---|--|
| Other   |  |
| Off-site<br>Burial                              | 22 L<br>26 L   |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 108,970<br>92,702<br>95,390<br>141,847<br>122,411<br>278,883 |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                         |
| Chemical  | Methyl<br>alcohol  |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

# Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the otner<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 4 L<br>21 L<br>14 L                                    |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     | •  |
| Product                                 | ·  |
| Quantity<br>Used<br>(kg)                | 575<br>689<br>1,628<br>1,949<br>3,009                  |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Methyl ethyl<br>ketone                                 |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Chemical<br>Name | Year | Quantity<br>Used<br>(kg) | Product | Airborne Liquid<br>Release Effluen | Liquid<br>Effluent | On-site<br>Storage | On-site Of<br>Burial B | ff-site<br>Burial Oth | On-site Off-site<br>Burial Burial Other Unknown |
|------------------|------|--------------------------|---------|------------------------------------|--------------------|--------------------|------------------------|-----------------------|---|
|                  |      |                          |         |                                    |                    |                    |                        |                       |   |
|                  |      | 5                        |         |                                    |                    |                    |                        |                       | Records do                                      |
| P,P'-Diamino-    | 1984 | 16                       |         |                                    |                    |                    |                        |                       | not exist for                                   |
| diphenylmethane  |      |                          |         |                                    |                    |                    |                        |                       | the other                                       |
|                  |      |                          |         |                                    |                    |                    |                        |                       | cateoories                                      |

## Estimated Distribution in %

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 4 L  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 9,470<br>9,746<br>13,017<br>15,333<br>9,637            |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Methylene<br>chloride                                  |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOOI). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 364<br>364<br>364<br>364<br>1,091                      |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Metex Stripper   |

| Chemical<br>Name | Year | Quantity<br>Used<br>(kg)                | Product | Airborne<br>Release | Liquid<br>Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Other | Unknown       |
|------------------|------|---|---------|---------------------|--------------------|--------------------|---|-------|---------------|
|                  |      |   | •       |                     |                    |                    |   |       |               |
|                  | 0    | ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, |         |                     |                    |                    |   |       | Records do    |
| Micro-Bio-Ireat  | 1983 | 7,074                                   |         |                     |                    |                    |   |       | not exist for |
|                  | 1982 | 2,082                                   |         |                     |                    |                    |   |       | for the other |
|                  | 1981 | 416                                     | -       |                     |                    |                    |   |       | categories    |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 4 L  |
| On-site On-<br>Storage Bur                      |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| y<br>Product                                    |  |
| Quantity<br>Used<br>(kg)                        | 6,245<br>4,788<br>6,037<br>3,331<br>2,498<br>1,665     |
| Year  | 1985<br>1984<br>1983<br>- 1982<br>1981                 |
| Chemical<br>Name                                | Naphtha  |

<sup>\*</sup>The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Chemical<br>Name | Year                                 | Quantity<br>Used<br>(kg)                   | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Ųnknown  |
|------------------|--------------------------------------|--|---------|-------------------------------------|--------------------|--------------------|---|--|
| .Nickel chloride | 1985<br>1984<br>1983<br>1982<br>1981 | 136<br>545<br>2,250<br>165<br>265<br>1,444 |         | -                                   | •                  |                    | < 1 kg<br>209 kg<br>17 kg               | Records do<br>not exist for<br>the other<br>categories |

| Unknown                                 | kecords do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | < 1 kg<br>< 1 kg<br>14 kg                              |
| On-site (<br>Storage E                  |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 173<br>236<br>227<br>827<br>1,127                      |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Nickel sulfate   |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories   |
|---|--|
| On-site Off-site<br>Burial Burial Other | 4 L<br>188 L<br>920 L                                    |
| On-site On-site<br>Storage Burial       | 106,832 L<br>Presently<br>in storage                     |
| Airborne Liquid<br>Release Effluent     | •  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 10,647<br>17,895<br>17,036<br>20,352<br>22,581<br>23,679 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                     |
| Chemical<br>Name                        | Nitric acid  |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      |  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 3<br>12<br>4   |
| Year                                    | 1985<br>1984<br>1983<br>1982                           |
| Chemical<br>Name                        | Nitric oxide   |

|                  | Unknown          | Records do<br>not exist for<br>the other<br>categories                       |
|------------------|------------------|--|
| On-site Off-site | al Burial Other  |  |
|                  |                  |  |
|                  | Storage          |  |
| Liauid           | Effluent         |  |
| Airhorne         | Release          |  |
| ٠                | Product          | · .  |
| Quantity         | (kg)             | 21-  |
|                  | Year             | 1983<br>1982<br>1981   |
|                  | Cnemical<br>Name | Orthene (Acetyl-<br>phosphoramido-<br>thioic acid-<br>0,5-dimethyl<br>ester) |

|  | Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|--|---|--|
|  | On-site Off-site<br>Burial Burial Other | 236 L<br>208 L   |
|  | On-site<br>Storage                      |  |
|  | Airborne Liquid<br>Release Effluent     |  |
|  | Airborne<br>Release                     |  |
|  | Product                                 |  |
|  | Quantity<br>Used<br>(kg)                | 15<br>42<br>19<br>4                                    |
|  | Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
|  | Chemical<br>Name                        | Pararosaniline.  |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | J L  |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             | -  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 30<br>38<br>76<br>238                                  |
| Year  | 1984<br>1983<br>1982<br>1981                           |
| Chemical<br>Name                                | Pentane  |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other |  |
| Off-s'<br>Buria                         | 12 kg<br>6 kg<br>34 kg                                 |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     | ·  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 494<br>290<br>279<br>358<br>306<br>291                 |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Perchloric acid  |

## Estimated Distribution in %

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | 16 L<br>24 L   |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             | ·  |
| Airborne<br>Release                             |  |
| Product   | ·  |
| Quantity<br>Used<br>(kg)                        | 193<br>189<br>238<br>314<br>193<br>379                 |
| Year  | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                                | Petroleum ether  |

\*The majority of solvent wastes occur as mixtures of several different chemicals and are managed as ignitable wastes (DOO1). These wastes are not mixed intentionally, rather, they are a result of the waste generating process.

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 9 kg<br>78 kg<br>56 kg                                 |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 | ·  |
| Quantity<br>Used<br>(kg)                | 29<br>40<br>22<br>40<br>77<br>35                       |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Phenol   |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 16   |
| Year  | 1985   |
| Chemical<br>Name                                | Phenolic resin   |

| Used Airborne<br>(kg) Product Release |
|---------------------------------------|
| ·                                     |

# Estimated Distribution in %

| Unknown   |                                     |
|---|-------------------------------------|
| -site<br>rial Other                             | 5,616 kg<br>18,000 kg<br>5,800 kg   |
| On-site Off-site<br>Burial Burial Other Unknown | 18,<br>5,                           |
| On-site<br>Storage                              |                                     |
| Liquid<br>Effluent                              |                                     |
| Airborne<br>Release 1                           |                                     |
| Product   | ·                                   |
| Quantity<br>Used<br>(kg)                        |                                     |
| Year  | 1985<br>1984<br>1983                |
| Chemical<br>Name                                | Polychlorinated<br>biphenyls (PCBs) |

\*There is approximately 25,200 L (> 500 ppm) and 3,017 L (50-500 ppm) PCB containing oils currently in use in various transformers and other electrical equipment.

|  | Unknown                                 | Records do<br>not exist for<br>the otner<br>categories |
|--|---|--|
|  | Other                                   |  |
|  | Off-site<br>Burial                      | 1 kg<br>2 kg<br>20 kg                                  |
|  | On-site Off-site<br>Burial Burial Other |  |
|  | On-site<br>Storage                      |  |
|  | Liquid<br>Effluent                      |  |
|  | Airborne Liquid<br>Release Effluent     |  |
|  | Product                                 | ·  |
|  | Quantity<br>Used<br>(kg)                | 88 68  |
|  | Year                                    | 1985<br>1984<br>1983<br>1981                           |
|  | Chemical<br>Name                        | Potassium<br>cyanide                                   |

| Chemical<br>Name       | Year                                 | Quantity<br>Used<br>(kg)                                   | Product | Airborne<br>Release  | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | Unknown  |
|------------------------|--------------------------------------|--|---------|----------------------|-------------------------------------|--------------------|-------------------|---|--|
| Potassium<br>hydroxide | 1985<br>1984<br>1983<br>1982<br>1981 | 113,220<br>157,750<br>60,597<br>48,674<br>92,850<br>84,586 | ·       |                      |                                     |                    |                   | 8 kg<br>27 kg<br>28 kg                  | Records do<br>not exist for<br>the other<br>categories |
|                        | 1970-1985                            | 8,180/yr.  | Used    | for cleaning<br>bers | <u>6</u> 1                          |                    |                   |   |  |

| her Unknown                             | Records do | not exist for | the other | categories |
|---|------------|---------------|-----------|------------|
| On-site Off-site<br>Burial Burial Other |            |               |           |            |
| On-site<br>Storage                      |            |               |           |            |
| Liquid<br>Effluent                      |            |               |           |            |
| Airborne<br>Release                     |            |               |           |            |
| Product                                 |            |               |           |            |
| Quantity<br>Used<br>(kg)                | 1          | 227           |           |            |
| Year                                    |            | 1983          |           |            |
| Chemical<br>Name                        |            | Pramitol      |           |            |

| ner Unknown                             | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other | 13 F<br>7 L<br>17 L                                    |
| On-sit<br>Burial                        |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 72 87  |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Pyridine   |

ESTIMATED CHEMICAL USAGE

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 455  |
| Year  | 1983   |
| Chemical<br>Name                                | Ronstar  |

| Chemical<br>Name   | Year                 | Quantity<br>Used<br>(kg) | Product | Airborne Liquid<br>Release Effluent | Liquid<br>Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Unknown  |
|--|----------------------|--------------------------|---------|-------------------------------------|--------------------|--------------------|---|--|
| Round-up<br>(N-phosphono-<br>methyl glycine<br>isopropylamine<br>salt) | 1984<br>1983<br>1982 | 11<br>182<br>38          | ·       |                                     |                    |                    |   | Records do<br>not exist for<br>the other<br>categories |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      |  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 3,125  |
| Year                                    | 1982   |
| Chemical<br>Name                        | Silicon<br>tetraflouride                               |

| Unknown   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              |  |
| On-site Off-site<br>Burial Burial Other Unknown |  |
| On-site O<br>Storage B                          |  |
| Liquid<br>Effluent                              |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(kg) F                      | 284  |
| Year  | 1983   |
| Chemical<br>Name                                | Silvex   |

| Unknown   | Records do<br>not exist for<br>the otner<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other Unknown | •  |
| On-site<br>Storage                              |  |
| Airborne Liquid<br>Release Effluent             |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 955<br>1,510<br>1,318<br>1,091                         |
| Year  | 1985<br>1983<br>1982<br>1981<br>1980                   |
| Chemical<br>Name                                | Sodium<br>bifluoride                                   |

| Unknown                                 | Records do<br>not exist for<br>the otner<br>categories |
|---|--|
| Other                                   |  |
| Off-site<br>Burial                      | 3 K G<br>3 K G<br>8 K G                                |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 92<br>2<br>1,002<br>365<br>182<br>727                  |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981                   |
| Chemical<br>Name                        | Sodium<br>cyanide                                      |

| Unknown   | Records do<br>not exist for<br>the other<br>categories       |
|---|--|
| Other   |  |
| On-site Off-site<br>Burial Burial Other Unknown | <ul><li> &gt; 1</li><li> &gt; 1</li><li> Y &amp; 9</li></ul> |
| On-site<br>Burial                               | ·  |
| On-site<br>Storage                              |  |
| e Liquid<br>Effluent                            |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 227<br>455<br>114<br>12                                      |
| Year  | 1985<br>1984<br>1983<br>1982                                 |
| Chemical<br>Name                                | Sodium<br>dithionite   |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                   | -  |
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Burial                       |  |
| On-site<br>Storage                      |  |
| Liquid<br>Effluent                      |  |
| Airborne Liquid<br>Release Effluen      |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg) P              | 455  |
| Year                                    | 1981   |
| Chemical<br>Name                        | Sodium<br>hexameta-<br>phosphate                       |

| Chemical<br>Name    | Year  | Quantity<br>Used<br>(kg)                                  | Product      | Airborne Liquid<br>Release Effluen | Liquid<br>Effluent | On-site<br>Storage                   | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | r Unknown  |
|---------------------|---|---|--------------|------------------------------------|--------------------|--------------------------------------|-------------------|---|--|
| Sodium<br>hydroxide | 1985<br>1984<br>1983<br>1982<br>1981<br>1980<br>1943-1979 | 106,138<br>97,897<br>75,132<br>53,889<br>35,462<br>26,129 | Steam Operat | Plant<br>ions                      | ·                  | 64, 534 L<br>Presently<br>in storage |                   | 19 kg<br>75 kg<br>48 kg                 | Records do<br>not exist for<br>the other<br>categories |

| Chemical<br>Name  | Year                                 | Quantity<br>Used<br>(kg)                               | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site<br>Burial | On-site Off-site<br>Burial Burial Other | Unknown  |
|-------------------|--------------------------------------|--|---------|---------------------|-------------------------------------|--------------------|-------------------|---|--|
| Sodium<br>nitrate | 1985<br>1984<br>1983<br>1982<br>1981 | 17,070<br>9,815<br>13,883<br>10,248<br>9,025<br>10,215 |         |                     |                                     |                    |                   | 2 kg<br>53 kg<br>3 kg                   | Records do<br>not exist for<br>the other<br>categories |

| Chemical<br>Name  | Year                 | Quantity<br>Used<br>(kg) | Product | Airborne<br>Release | Airborne Liquid<br>Release Effluent | On-site<br>Storage | On-site Off-site<br>Burial Burial Other | Unknown  |
|-------------------|----------------------|--------------------------|---------|---------------------|-------------------------------------|--------------------|---|--|
| Sodium<br>sulfide | 1985<br>1984<br>1983 | 200                      | -       |                     |                                     |                    | 3 kg<br>4 kg<br>3 kg                    | Records do<br>not exist for<br>the other<br>categories |

| er Unknown                              | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| On-site Off-site<br>Burial Burial Other |  |
| On-site<br>Storage                      |  |
| Airborne Liquid<br>Release Effluent     |  |
| Airborne<br>Release                     |  |
| Product                                 |  |
| Quantity<br>Used<br>(kg)                | 4  |
| Year                                    | 1982   |
| Chemical<br>Name                        | Subdue E2  |

| Unknown                                   | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other                                     |  |
| On-site Off-site<br>Burial Burial Other L | r  |
| On-Site<br>Burial                         |  |
| On-site<br>Storage                        |  |
| Liquid<br>Effluent                        |  |
| Airborne Liquid<br>Release Effluent       |  |
| Product                                   |  |
| Quantity<br>Used<br>(kg)                  | 7,214  |
| Year                                      | 1981   |
| Chemical<br>Name                          | Sulfur<br>hexafluoride                                 |

| Unknown                                 | Records do<br>not exist for<br>the other<br>categories          |
|---|---|
| On-site Off-site<br>Burial Burial Other | 7 L<br>75 L<br>81 L   |
| On-site On-site<br>Storage Burial       | 41,143 L<br>Presently<br>in storage                             |
|   | 41,<br>Pres   |
| Airborne Liquid<br>Release Effluent     | Plant<br>tions  |
| Product                                 | Steam<br>Operat   |
| Quantity<br>Used<br>(kg)                | 7,305<br>9,606<br>11,582<br>5,185<br>5,042<br>7,244<br>3,800/yr |
| Year                                    | 1985<br>1984<br>1983<br>1982<br>1981<br>1981<br>1980            |
| Chemical<br>Name                        | Sulfuric  |

C

| On-site Off-site<br>Burial Burial Other Unknown | Records do<br>not exist for<br>the other<br>categories |
|---|--|
| Other   |  |
| Off-site<br>Burial                              |  |
| On-site<br>Burial                               |  |
| On-site<br>Storage                              |  |
| Liquid<br>Effluent                              |  |
| Airborne<br>Release                             |  |
| Product   |  |
| Quantity<br>Used<br>(kg)                        | 23<br>23<br>23   |
| Year  | 1984<br>1982<br>1981                                   |
| Chemical<br>Name                                | Talon (4'-Bromo<br>(1,1'-biphenyl-<br>4-yl)            |